

Table 12.14
Unadjusted Patient Survival, Deceased Donor Lung Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

| | 3 Months | | | 1 Year | | | 5 Years | | | 10 Years | | |
|---------------------------------|------------------|--------|-----------|------------------|-------|-----------|------------------|--------|-----------|------------------|-------|-----------|
| | (Tx 2007 - 2008) | | | (Tx 2007 - 2008) | | | (Tx 2003 - 2008) | | | (Tx 1998 - 2008) | | |
| | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. |
| Total | | | | | | | | | | | | |
| All | 2,777 | 92.5% | 0.5% | 2,777 | 83.0% | 0.7% | 7,599 | 54.1% | 0.7% | 12,189 | 28.8% | 0.7% |
| Age at Tx | | | | | | | | | | | | |
| <1 Year | 7 | 100.0% | 0.0% | 7 | 85.7% | 13.2% | 22 | 44.3% | 13.6% | 40 | 40.3% | 9.4% |
| 1-5 Years | 8 | 75.0% | 15.3% | 8 | 75.0% | 15.3% | 23 | 60.9% | 11.7% | 46 | 40.5% | 9.4% |
| 6-11 Years | 10 | 100.0% | 0.0% | 10 | 90.0% | 9.5% | 48 | 52.0% | 8.3% | 92 | 37.5% | 7.0% |
| 12-17 Years | 63 | 96.8% | 2.2% | 63 | 88.9% | 4.0% | 178 | 50.7% | 4.7% | 272 | 27.1% | 4.3% |
| 18-34 Years | 304 | 96.7% | 1.0% | 304 | 86.8% | 1.9% | 898 | 52.2% | 2.1% | 1,489 | 37.7% | 1.8% |
| 35-49 Years | 398 | 94.2% | 1.2% | 398 | 87.9% | 1.6% | 1,245 | 61.0% | 1.7% | 2,316 | 37.1% | 1.5% |
| 50-64 Years | 1,508 | 91.6% | 0.7% | 1,508 | 82.4% | 1.0% | 4,263 | 54.9% | 1.0% | 6,831 | 24.6% | 1.0% |
| 65+ Years | 479 | 91.0% | 1.3% | 479 | 77.5% | 1.9% | 922 | 41.7% | 2.4% | 1,103 | 10.8% | 2.7% |
| Recipient Ethnicity/Race | | | | | | | | | | | | |
| White | 2,315 | 92.5% | 0.5% | 2,315 | 82.6% | 0.8% | 6,466 | 54.2% | 0.8% | 10,576 | 29.0% | 0.7% |
| African American | 245 | 93.5% | 1.6% | 245 | 86.9% | 2.2% | 622 | 55.0% | 2.6% | 922 | 25.5% | 3.0% |
| Hispanic/Latino | 160 | 92.5% | 2.1% | 160 | 83.1% | 3.0% | 370 | 51.1% | 3.4% | 504 | 29.6% | 3.8% |
| Asian | 37 | 91.9% | 4.5% | 37 | 81.1% | 6.4% | 92 | 55.1% | 6.7% | 121 | 44.3% | 6.2% |
| Other/Multi-race | 20 | 85.0% | 8.0% | 20 | 80.0% | 8.9% | 47 | 43.7% | 10.6% | 64 | 24.2% | 11.2% |
| Unknown | 0 | - | - | 0 | - | - | 2 | 100.0% | 0.0% | 2 | 0.0% | 0.0% |
| Recipient Gender | | | | | | | | | | | | |
| Female | 1,126 | 93.5% | 0.7% | 1,126 | 84.4% | 1.1% | 3,339 | 54.8% | 1.1% | 5,645 | 30.2% | 1.0% |
| Male | 1,651 | 91.9% | 0.7% | 1,651 | 82.1% | 0.9% | 4,260 | 53.5% | 1.0% | 6,544 | 27.5% | 1.0% |
| Recipient Blood Type | | | | | | | | | | | | |
| O | 1,247 | 93.0% | 0.7% | 1,247 | 84.0% | 1.0% | 3,405 | 53.8% | 1.1% | 5,364 | 28.3% | 1.1% |
| A | 1,113 | 92.5% | 0.8% | 1,113 | 83.5% | 1.1% | 3,073 | 54.4% | 1.1% | 4,965 | 29.5% | 1.1% |
| B | 292 | 90.8% | 1.7% | 292 | 78.4% | 2.4% | 814 | 54.1% | 2.2% | 1,360 | 28.8% | 2.1% |
| AB | 125 | 92.0% | 2.4% | 125 | 80.0% | 3.6% | 307 | 54.3% | 3.7% | 500 | 26.8% | 3.3% |

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on lung and heart-lung transplants.

Multi-organ transplants are excluded.

Table 12.14 (Continued)
Unadjusted Patient Survival, Deceased Donor Lung Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

| | 3 Months | | | 1 Year | | | 5 Years | | | 10 Years | | |
|--|------------------|--------|-----------|------------------|--------|-----------|------------------|-------|-----------|------------------|-------|-----------|
| | (Tx 2007 - 2008) | | | (Tx 2007 - 2008) | | | (Tx 2003 - 2008) | | | (Tx 1998 - 2008) | | |
| | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. |
| Total | | | | | | | | | | | | |
| All | 2,777 | 92.5% | 0.5% | 2,777 | 83.0% | 0.7% | 7,599 | 54.1% | 0.7% | 12,189 | 28.8% | 0.7% |
| Recipient Residency | | | | | | | | | | | | |
| U.S. Resident | 2,759 | 92.6% | 0.5% | 2,759 | 83.0% | 0.7% | 7,564 | 54.1% | 0.7% | 12,118 | 28.7% | 0.7% |
| Nonresident Alien | 18 | 88.9% | 7.4% | 18 | 83.3% | 8.8% | 33 | 67.1% | 9.6% | 58 | 54.7% | 7.6% |
| Unknown | 0 | - | - | 0 | - | - | 2 | 50.0% | 35.4% | 13 | 35.9% | 13.9% |
| Hospitalized at Tx | | | | | | | | | | | | |
| Not Hospitalized | 2,351 | 94.0% | 0.5% | 2,351 | 85.2% | 0.7% | 6,668 | 55.8% | 0.8% | 10,912 | 29.0% | 0.8% |
| Hospitalized | 209 | 92.3% | 1.8% | 209 | 79.9% | 2.8% | 489 | 42.0% | 3.4% | 694 | 30.4% | 2.9% |
| In Intensive Care | 217 | 77.4% | 2.8% | 217 | 62.7% | 3.3% | 442 | 40.1% | 2.8% | 572 | 24.6% | 3.2% |
| Unknown | 0 | - | - | 0 | - | - | 0 | - | - | 11 | 43.6% | 15.5% |
| Life Support at Tx | | | | | | | | | | | | |
| No | 2,579 | 93.4% | 0.5% | 2,579 | 83.9% | 0.7% | 7,135 | 54.5% | 0.7% | 11,475 | 28.9% | 0.7% |
| Yes | 198 | 81.8% | 2.7% | 198 | 71.2% | 3.2% | 464 | 48.3% | 2.8% | 703 | 28.4% | 3.1% |
| Unknown | 0 | - | - | 0 | - | - | 0 | - | - | 11 | 43.6% | 15.5% |
| Primary Diagnosis | | | | | | | | | | | | |
| Emphysema/COPD | 801 | 92.9% | 0.9% | 801 | 83.1% | 1.3% | 2,500 | 54.7% | 1.2% | 4,473 | 24.0% | 1.1% |
| Cystic Fibrosis | 373 | 96.5% | 0.9% | 373 | 87.1% | 1.7% | 1,120 | 56.3% | 1.8% | 1,850 | 38.8% | 1.7% |
| Idiopathic Pulmonary Fibrosis | 950 | 92.0% | 0.9% | 950 | 81.6% | 1.3% | 2,213 | 52.0% | 1.4% | 2,989 | 24.7% | 1.8% |
| Alpha-1-Antitrypsin Deficiency | 78 | 97.4% | 1.8% | 78 | 91.0% | 3.2% | 293 | 61.7% | 3.2% | 655 | 30.8% | 2.8% |
| Idiopathic Pulmonary Arterial Hypertension | 48 | 87.5% | 4.8% | 48 | 83.3% | 5.4% | 182 | 51.0% | 4.4% | 380 | 38.6% | 3.2% |
| Congenital Disease | 2 | 100.0% | 0.0% | 2 | 100.0% | 0.0% | 19 | 44.9% | 11.9% | 79 | 36.8% | 6.0% |
| Other | 524 | 89.9% | 1.3% | 524 | 81.1% | 1.7% | 1,271 | 53.7% | 1.9% | 1,762 | 33.8% | 2.0% |
| Unknown | 1 | 100.0% | 0.0% | 1 | 100.0% | 0.0% | 1 | + | + | 1 | + | + |

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on lung and heart-lung transplants.

Multi-organ transplants are excluded.

Table 12.14 (Continued)
Unadjusted Patient Survival, Deceased Donor Lung Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

| | 3 Months | | | 1 Year | | | 5 Years | | | 10 Years | | |
|-----------------------------|------------------|-------|-----------|------------------|-------|-----------|------------------|-------|-----------|------------------|-------|-----------|
| | (Tx 2007 - 2008) | | | (Tx 2007 - 2008) | | | (Tx 2003 - 2008) | | | (Tx 1998 - 2008) | | |
| | N | % | Std. Err. |
| Total | | | | | | | | | | | | |
| All | 2,777 | 92.5% | 0.5% | 2,777 | 83.0% | 0.7% | 7,599 | 54.1% | 0.7% | 12,189 | 28.8% | 0.7% |
| Cold Ischemic Time | | | | | | | | | | | | |
| 0-90 Minutes | 11 | 81.8% | 11.6% | 11 | 72.7% | 13.4% | 46 | 45.3% | 8.2% | 97 | 17.7% | 5.7% |
| 91-180 Minutes | 191 | 93.7% | 1.8% | 191 | 85.9% | 2.5% | 658 | 51.0% | 2.4% | 1,327 | 22.1% | 1.7% |
| 181-270 Minutes | 754 | 92.4% | 1.0% | 754 | 83.0% | 1.4% | 2,097 | 51.3% | 1.4% | 3,477 | 24.8% | 1.3% |
| 271-360 Minutes | 954 | 92.0% | 0.9% | 954 | 82.4% | 1.2% | 2,443 | 56.3% | 1.3% | 3,603 | 34.0% | 1.4% |
| 361-480 Minutes | 545 | 93.4% | 1.1% | 545 | 84.6% | 1.5% | 1,337 | 55.9% | 1.7% | 1,806 | 34.3% | 2.3% |
| 481+ Minutes | 140 | 90.7% | 2.5% | 140 | 77.1% | 3.5% | 313 | 55.3% | 3.7% | 465 | 38.8% | 3.7% |
| Unknown | 182 | 94.0% | 1.8% | 182 | 83.5% | 2.8% | 705 | 55.1% | 2.3% | 1,414 | 28.4% | 1.9% |
| Yearly Center Volume | | | | | | | | | | | | |
| 0-3 | 39 | 89.7% | 4.9% | 39 | 82.1% | 6.1% | 145 | 62.5% | 5.2% | 506 | 26.8% | 2.7% |
| 4-6 | 35 | 88.6% | 5.4% | 35 | 88.6% | 5.4% | 232 | 43.7% | 4.0% | 165 | 28.3% | 4.8% |
| 7-11 | 209 | 94.3% | 1.6% | 209 | 82.3% | 2.6% | 272 | 50.5% | 3.9% | 1,096 | 22.5% | 2.5% |
| 12-20 | 393 | 90.8% | 1.5% | 393 | 79.1% | 2.0% | 1,118 | 46.4% | 1.9% | 2,733 | 24.2% | 1.4% |
| 21+ | 2,101 | 92.8% | 0.6% | 2,101 | 83.7% | 0.8% | 5,832 | 56.0% | 0.8% | 7,689 | 31.3% | 0.9% |
| Type of Procedure | | | | | | | | | | | | |
| Double Lung | 1,833 | 92.3% | 0.6% | 1,833 | 83.6% | 0.9% | 4,655 | 56.7% | 0.9% | 6,734 | 37.0% | 1.0% |
| Single Lung | 944 | 93.0% | 0.8% | 944 | 81.9% | 1.3% | 2,944 | 50.5% | 1.1% | 5,455 | 21.6% | 0.9% |
| Donor Age | | | | | | | | | | | | |
| <1 Year | 9 | 88.9% | 10.5% | 9 | 77.8% | 13.9% | 23 | 46.6% | 12.2% | 40 | 37.2% | 9.8% |
| 1-5 Years | 8 | 87.5% | 11.7% | 8 | 75.0% | 15.3% | 38 | 49.0% | 10.2% | 80 | 30.6% | 6.5% |
| 6-11 Years | 30 | 96.7% | 3.3% | 30 | 86.7% | 6.2% | 113 | 52.2% | 5.4% | 218 | 28.8% | 4.9% |
| 12-17 Years | 308 | 92.5% | 1.5% | 308 | 82.1% | 2.2% | 943 | 52.8% | 2.0% | 1,613 | 32.0% | 1.7% |
| 18-34 Years | 1,242 | 92.7% | 0.7% | 1,242 | 84.0% | 1.0% | 3,320 | 54.4% | 1.1% | 5,220 | 30.7% | 1.1% |
| 35-49 Years | 713 | 93.1% | 0.9% | 713 | 83.2% | 1.4% | 1,944 | 55.8% | 1.4% | 3,250 | 25.8% | 1.3% |
| 50-64 Years | 441 | 91.8% | 1.3% | 441 | 82.1% | 1.8% | 1,142 | 53.4% | 1.9% | 1,679 | 25.5% | 2.2% |
| 65+ Years | 26 | 80.8% | 7.7% | 26 | 57.7% | 9.7% | 76 | 32.5% | 6.4% | 89 | 19.8% | 9.1% |

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on lung and heart-lung transplants.

Multi-organ transplants are excluded.

Table 12.14 (Continued)
Unadjusted Patient Survival, Deceased Donor Lung Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

| | 3 Months | | | 1 Year | | | 5 Years | | | 10 Years | | |
|--|------------------|--------|-----------|------------------|--------|-----------|------------------|-------|-----------|------------------|-------|-----------|
| | (Tx 2007 - 2008) | | | (Tx 2007 - 2008) | | | (Tx 2003 - 2008) | | | (Tx 1998 - 2008) | | |
| | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. |
| Total | | | | | | | | | | | | |
| All | 2,777 | 92.5% | 0.5% | 2,777 | 83.0% | 0.7% | 7,599 | 54.1% | 0.7% | 12,189 | 28.8% | 0.7% |
| Lung Diagnosis Group | | | | | | | | | | | | |
| Group A: Obstructive Lung Disease | 989 | 93.2% | 0.8% | 989 | 84.0% | 1.2% | 3,151 | 55.4% | 1.1% | 5,706 | 25.8% | 1.0% |
| Group B: Pulmonary Vascular Disease | 115 | 85.2% | 3.3% | 115 | 80.0% | 3.7% | 311 | 50.0% | 3.7% | 541 | 38.5% | 2.9% |
| Group C: Cystic Fibrosis or Immunodef. Disorders | 372 | 96.0% | 1.0% | 372 | 86.6% | 1.8% | 1,101 | 56.8% | 1.8% | 1,808 | 39.0% | 1.7% |
| Group D: Restrictive Lung Disease | 1,232 | 91.7% | 0.8% | 1,232 | 81.2% | 1.1% | 2,861 | 52.1% | 1.3% | 3,825 | 26.2% | 1.6% |
| Group E: Age Less Than 12 | 25 | 92.0% | 5.4% | 25 | 84.0% | 7.3% | 93 | 51.7% | 6.2% | 178 | 38.5% | 4.9% |
| Other Diagnosis | 44 | 90.9% | 4.3% | 44 | 88.6% | 4.8% | 82 | 59.6% | 8.2% | 131 | 38.7% | 7.0% |
| Lung Allocation Score | | | | | | | | | | | | |
| LAS 0 | 2 | 100.0% | 0.0% | 2 | 100.0% | 0.0% | 0 | - | - | 0 | - | - |
| LAS 20 - 29 | 19 | 94.7% | 5.1% | 19 | 89.5% | 7.0% | 0 | - | - | 0 | - | - |
| LAS 30 - 34 | 804 | 93.9% | 0.8% | 804 | 85.9% | 1.2% | 0 | - | - | 0 | - | - |
| LAS 35 - 39 | 698 | 94.1% | 0.9% | 698 | 82.8% | 1.4% | 0 | - | - | 0 | - | - |
| LAS 40 - 49 | 668 | 91.5% | 1.1% | 668 | 82.6% | 1.5% | 0 | - | - | 0 | - | - |
| LAS 50 - 59 | 225 | 92.0% | 1.8% | 225 | 82.2% | 2.5% | 0 | - | - | 0 | - | - |
| LAS 60+ | 361 | 88.6% | 1.7% | 361 | 77.6% | 2.2% | 0 | - | - | 0 | - | - |
| Unknown | 0 | - | - | 0 | - | - | 7,599 | 54.1% | 0.7% | 12,189 | 28.8% | 0.7% |

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on lung and heart-lung transplants.

Multi-organ transplants are excluded.

Table 12.14 (Continued)
Unadjusted Patient Survival, Deceased Donor Lung Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

| | 3 Months | | | 1 Year | | | 5 Years | | | 10 Years | | |
|--------------------------------|------------------|--------|-----------|------------------|--------|-----------|------------------|-------|-----------|------------------|-------|-----------|
| | (Tx 2007 - 2008) | | | (Tx 2007 - 2008) | | | (Tx 2003 - 2008) | | | (Tx 1998 - 2008) | | |
| | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. |
| Total | | | | | | | | | | | | |
| All | 2,777 | 92.5% | 0.5% | 2,777 | 83.0% | 0.7% | 7,599 | 54.1% | 0.7% | 12,189 | 28.8% | 0.7% |
| Transplant Center State | | | | | | | | | | | | |
| Alabama | 53 | 90.6% | 4.0% | 53 | 81.1% | 5.4% | 164 | 50.4% | 4.8% | 324 | 26.0% | 3.5% |
| Arizona | 53 | 90.6% | 4.0% | 53 | 84.9% | 4.9% | 87 | 52.3% | 8.3% | 113 | 37.5% | 8.5% |
| California | 361 | 94.2% | 1.2% | 361 | 83.9% | 1.9% | 978 | 52.0% | 2.0% | 1,431 | 23.6% | 2.4% |
| Colorado | 60 | 96.7% | 2.3% | 60 | 86.7% | 4.4% | 190 | 49.2% | 4.3% | 338 | 23.5% | 3.8% |
| Florida | 205 | 91.7% | 1.9% | 205 | 84.4% | 2.5% | 535 | 59.7% | 2.7% | 699 | 31.3% | 3.8% |
| Georgia | 49 | 95.9% | 2.8% | 49 | 89.8% | 4.3% | 145 | 52.0% | 5.3% | 199 | 20.2% | 5.2% |
| Iowa | 17 | 100.0% | 0.0% | 17 | 94.1% | 5.7% | 17 | + | + | 17 | + | + |
| Illinois | 102 | 93.1% | 2.5% | 102 | 83.3% | 3.7% | 234 | 64.6% | 3.7% | 395 | 34.1% | 3.9% |
| Indiana | 56 | 85.7% | 4.7% | 56 | 71.4% | 6.0% | 224 | 50.7% | 3.9% | 349 | 18.0% | 3.8% |
| Kentucky | 58 | 89.7% | 4.0% | 58 | 70.7% | 6.0% | 141 | 45.5% | 5.4% | 240 | 19.7% | 4.0% |
| Louisiana | 0 | - | - | 0 | - | - | 39 | 56.4% | 7.9% | 127 | 10.0% | 3.8% |
| Massachusetts | 67 | 97.0% | 2.1% | 67 | 85.1% | 4.4% | 190 | 42.3% | 4.5% | 332 | 25.5% | 4.0% |
| Maryland | 62 | 91.9% | 3.5% | 62 | 83.9% | 4.7% | 157 | 43.8% | 5.9% | 295 | 30.1% | 3.9% |
| Michigan | 77 | 93.5% | 2.8% | 77 | 85.7% | 4.0% | 205 | 60.9% | 4.4% | 397 | 34.4% | 3.8% |
| Minnesota | 85 | 94.1% | 2.6% | 85 | 82.4% | 4.1% | 272 | 62.4% | 3.7% | 452 | 42.3% | 3.4% |
| Missouri | 129 | 93.8% | 2.1% | 129 | 83.7% | 3.3% | 380 | 62.8% | 3.0% | 721 | 42.7% | 2.6% |
| North Carolina | 143 | 92.3% | 2.2% | 143 | 86.0% | 2.9% | 421 | 53.6% | 3.1% | 733 | 30.8% | 2.8% |
| Nebraska | 0 | - | - | 0 | - | - | 0 | - | - | 6 | 16.7% | 15.2% |
| New Jersey | 3 | 100.0% | 0.0% | 3 | 100.0% | 0.0% | 3 | + | + | 3 | + | + |
| New York | 108 | 96.3% | 1.8% | 108 | 86.1% | 3.3% | 287 | 59.1% | 3.8% | 410 | 33.0% | 4.5% |
| Ohio | 142 | 90.8% | 2.4% | 142 | 84.5% | 3.0% | 408 | 55.9% | 3.0% | 668 | 25.6% | 2.9% |
| Oklahoma | 22 | 95.5% | 4.4% | 22 | 77.3% | 8.9% | 83 | 29.4% | 6.3% | 122 | 19.5% | 5.5% |
| Oregon | 0 | - | - | 0 | - | - | 0 | - | - | 6 | 33.3% | 19.2% |
| Pennsylvania | 403 | 89.8% | 1.5% | 403 | 78.7% | 2.0% | 900 | 55.9% | 2.1% | 1,327 | 28.8% | 2.4% |
| Tennessee | 28 | 92.9% | 4.9% | 28 | 85.7% | 6.6% | 116 | 36.7% | 5.4% | 203 | 27.0% | 3.9% |
| Texas | 226 | 90.3% | 2.0% | 226 | 83.6% | 2.5% | 592 | 49.2% | 2.6% | 917 | 25.4% | 2.5% |
| Utah | 18 | 100.0% | 0.0% | 18 | 100.0% | 0.0% | 46 | 53.7% | 9.3% | 88 | 31.0% | 6.7% |
| Virginia | 80 | 91.3% | 3.2% | 80 | 77.5% | 4.7% | 296 | 52.0% | 3.6% | 497 | 22.6% | 3.3% |
| Washington | 93 | 96.8% | 1.8% | 93 | 90.3% | 3.1% | 267 | 61.1% | 3.9% | 395 | 22.7% | 6.7% |
| Wisconsin | 77 | 93.5% | 2.8% | 77 | 77.9% | 4.7% | 222 | 58.3% | 4.0% | 385 | 42.5% | 3.6% |

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on lung and heart-lung transplants.

Multi-organ transplants are excluded.