

Table 13.12
Adjusted Patient Survival, Heart-Lung Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2007 - 2008)			(Tx 2007 - 2008)			(Tx 2003 - 2008)			(Tx 1998 - 2008)		
	N	%	Std. Err.									
Total												
All	57	86.4%	4.5%	57	79.5%	5.4%	186	49.7%	4.1%	383	35.4%	3.1%
Age at Tx												
<1 Year	1	+	+	1	+	+	4	25.0%	16.3%	11	9.1%	6.8%
1-5 Years	3	66.7%	22.2%	3	66.7%	22.2%	6	55.6%	20.7%	14	38.1%	12.9%
6-11 Years	2	+	+	2	+	+	6	66.7%	17.4%	12	58.3%	13.5%
12-17 Years	3	+	+	3	+	+	15	48.0%	13.0%	30	42.0%	10.2%
18-34 Years	17	82.4%	8.9%	17	70.6%	10.7%	51	46.2%	7.5%	102	34.8%	5.6%
35-49 Years	21	85.7%	7.4%	21	81.0%	8.3%	73	48.8%	6.4%	144	35.1%	5.2%
50-64 Years	10	90.0%	9.0%	10	80.0%	12.0%	31	59.1%	9.8%	70	35.9%	7.0%
Recipient Ethnicity/Race												
White	40	90.2%	4.6%	40	80.4%	6.3%	130	50.1%	4.9%	296	35.6%	3.5%
African American	5	81.6%	15.3%	5	81.6%	15.3%	19	57.2%	11.1%	31	40.0%	10.4%
Hispanic/Latino	7	58.1%	17.6%	7	58.1%	17.6%	22	53.9%	10.6%	36	37.3%	9.8%
Asian	4	+	+	4	+	+	13	27.7%	14.0%	14	26.0%	13.2%
Other/Multi-race	1	+	+	1	+	+	2	0.0%	0.0%	6	20.7%	14.0%
Recipient Gender												
Female	37	92.2%	4.3%	37	84.4%	5.9%	105	48.0%	5.6%	220	38.5%	4.3%
Male	20	75.4%	9.4%	20	70.4%	10.0%	81	51.9%	5.8%	163	30.7%	4.5%

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on heart-lung transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Table 13.12 (Continued)
Adjusted Patient Survival, Heart-Lung Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2007 - 2008)			(Tx 2007 - 2008)			(Tx 2003 - 2008)			(Tx 1998 - 2008)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	57	86.4%	4.5%	57	79.5%	5.4%	186	49.7%	4.1%	383	35.4%	3.1%
Primary Diagnosis												
Congenital Disease	6	84.5%	13.4%	6	68.8%	17.4%	40	47.7%	8.2%	115	27.1%	5.2%
Cystic Fibrosis	2	+	+	2	+	+	6	44.4%	21.8%	9	22.3%	14.3%
Emphysema/COPD	0	-	-	0	-	-	2	0.0%	0.0%	12	38.9%	13.9%
Other	9	100.0%	0.0%	9	89.8%	9.4%	28	56.5%	10.9%	39	41.5%	8.6%
Unknown	24	79.5%	8.1%	24	71.3%	9.1%	59	54.9%	6.7%	111	39.7%	5.4%

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on heart-lung transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Table 13.12 (Continued)
Adjusted Patient Survival, Heart-Lung Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2007 - 2008)			(Tx 2007 - 2008)			(Tx 2003 - 2008)			(Tx 1998 - 2008)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	57	86.4%	4.5%	57	79.5%	5.4%	186	49.7%	4.1%	383	35.4%	3.1%
Transplant Center State												
Alabama	1	+	+	1	+	+	3	34.8%	20.4%	14	31.1%	12.3%
Arizona	1	0.0%	0.0%	1	0.0%	0.0%	5	22.1%	14.3%	15	47.1%	12.3%
California	24	91.8%	5.5%	24	87.7%	6.5%	49	48.2%	8.6%	87	26.3%	7.6%
Colorado	0	-	-	0	-	-	2	52.2%	24.6%	5	30.3%	17.9%
Florida	2	+	+	2	+	+	13	69.7%	12.2%	18	26.5%	15.1%
Illinois	0	-	-	0	-	-	4	52.2%	20.6%	7	56.3%	17.1%
Massachusetts	0	-	-	0	-	-	0	-	-	1	0.0%	0.0%
Maryland	4	78.3%	18.0%	4	78.3%	18.0%	7	57.3%	17.2%	10	49.5%	14.6%
Michigan	0	-	-	0	-	-	0	-	-	2	50.3%	25.0%
Minnesota	1	+	+	1	+	+	6	52.0%	17.9%	18	42.3%	11.5%
Missouri	1	+	+	1	+	+	5	58.1%	20.2%	17	25.5%	11.7%
North Carolina	1	+	+	1	+	+	9	35.6%	14.4%	25	31.5%	9.1%
New York	0	-	-	0	-	-	3	65.4%	22.4%	15	13.1%	8.4%
Ohio	5	+	+	5	+	+	18	51.3%	11.4%	20	44.7%	12.3%
Oregon	0	-	-	0	-	-	0	-	-	1	0.0%	0.0%
Pennsylvania	13	75.4%	11.7%	13	58.7%	13.4%	42	41.6%	8.7%	82	36.8%	5.9%
Tennessee	0	-	-	0	-	-	3	67.6%	21.7%	9	22.7%	13.6%
Texas	2	42.0%	22.6%	2	42.0%	22.6%	12	65.7%	13.2%	21	53.4%	11.3%
Utah	0	-	-	0	-	-	0	-	-	1	0.0%	0.0%
Washington	1	+	+	1	+	+	1	+	+	3	+	+
Wisconsin	1	100.0%	0.0%	1	0.0%	0.0%	4	72.8%	19.7%	12	49.4%	13.7%

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on heart-lung transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.