

**Table 5.13b**  
**Adjusted Patient Survival by Year of Transplant at 3 Months, 1 Year,**  
**3 Years, 5 Years and 10 Years**  
**Deceased Donor ECD Kidney Transplants**

Trans- # plant Year	3 Months		1 Year		3 Years		5 Years		10 Years	
	Surv.	Std. Err.	Surv.	Std. Err.	Surv.	Std. Err.	Surv.	Std. Err.	Surv.	Std. Err.
Trans- # plant Year										
1987 37	82.8%	6.3%	73.9%	7.3%	73.9%	7.3%	61.5%	8.2%	35.6%	8.3%
1988 149	96.7%	1.6%	91.8%	2.5%	70.9%	4.1%	52.6%	4.4%	33.2%	4.1%
1989 192	95.2%	1.6%	88.7%	2.4%	79.7%	3.1%	67.0%	3.6%	41.6%	3.8%
1990 286	95.2%	1.3%	86.5%	2.1%	76.2%	2.7%	68.6%	2.9%	40.2%	3.1%
1991 348	95.9%	1.1%	91.7%	1.6%	81.2%	2.2%	69.1%	2.6%	39.6%	2.8%
1992 434	95.4%	1.1%	90.6%	1.5%	79.0%	2.1%	67.5%	2.4%	39.0%	2.5%
1993 471	94.0%	1.1%	88.6%	1.5%	78.1%	2.0%	64.7%	2.3%	40.4%	2.4%
1994 738	97.6%	0.6%	92.4%	1.0%	82.6%	1.4%	71.3%	1.7%	43.0%	1.9%
1995 883	96.0%	0.7%	91.8%	0.9%	82.6%	1.3%	71.2%	1.6%	43.5%	1.8%
1996 971	96.3%	0.6%	92.0%	0.9%	83.3%	1.2%	72.3%	1.4%	46.4%	1.6%
1997 1,020	95.6%	0.6%	89.5%	0.9%	79.8%	1.2%	69.1%	1.4%	43.5%	1.6%
1998 1,084	97.0%	0.5%	90.9%	0.8%	80.3%	1.2%	70.6%	1.4%	45.1%	1.5%
1999 1,101	95.8%	0.6%	91.4%	0.8%	82.4%	1.1%	71.7%	1.3%	48.2%	1.5%
2000 1,044	96.4%	0.5%	91.0%	0.8%	82.8%	1.1%	71.3%	1.4%	+	+
2001 1,084	96.2%	0.5%	90.2%	0.9%	81.5%	1.1%	69.5%	1.4%	+	+
2002 1,184	96.0%	0.5%	92.1%	0.7%	83.7%	1.0%	74.2%	1.2%	+	+
2003 1,324	96.9%	0.4%	91.8%	0.7%	83.4%	0.9%	75.1%	1.1%	+	+
2004 1,399	97.2%	0.4%	92.3%	0.6%	82.9%	0.9%	73.5%	1.1%	+	+
2005 1,592	97.3%	0.4%	92.2%	0.6%	85.0%	0.8%	76.3%	1.2%	+	+
2006 1,624	97.0%	0.4%	93.1%	0.6%	85.9%	0.8%	+	+	+	+
2007 1,666	98.1%	0.3%	94.8%	0.5%	+	+	+	+	+	+
2008 1,768	98.2%	0.3%	94.8%	0.5%	+	+	+	+	+	+

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up.

Multi-organ transplants are excluded for kidney, pancreas, heart, lung, and liver survival rates.

Intestine survival includes intestine only and intestine with liver and/or pancreas transplants.

Living donor transplants excluded unless explicitly listed. Heterotopic heart and liver transplants excluded.

Patient survival follows patients from first transplant of this type until death. Graft survival follows individual transplants until graft failure. Counts for patient and graft survival are different because a patient may have more than one transplant for a type organ.

Survival rates are adjusted to the characteristics of transplants in year 1999. See Technical Notes for details of patient and graft survival methods.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.