

Table 7.6e
Immunosuppression Use for Maintenance
Prior to Discharge, 2000 to 2009
Recipients with Pancreas After Kidney (PAK) Transplants

	Year of Transplant									
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Functioning Graft at Discharge	278	287	336	315	380	306	255	240	192	183
With Immunosuppression Info	277	286	335	312	375	303	249	230	191	178
Corticosteroids										
Any in Category	95.7%	93.4%	89.9%	80.8%	74.7%	66.7%	63.9%	57.8%	60.2%	59.6%
Steroids	95.7%	93.4%	89.9%	80.8%	74.7%	66.7%	63.9%	57.8%	60.2%	59.6%
Cyclosporine										
Any in Category	13.7%	9.8%	11.3%	8.3%	8.3%	6.6%	5.6%	4.8%	3.7%	3.9%
Sandimmune	2.2%	0.7%	0.6%	0.3%	0.5%	0.3%	0.0%	1.7%	0.0%	0.0%
Neoral	11.6%	8.0%	9.0%	6.1%	5.6%	5.9%	5.2%	3.0%	3.1%	2.8%
Gengraf	0.0%	1.0%	1.8%	1.9%	2.1%	0.3%	0.4%	0.0%	0.5%	1.1%
Tacrolimus										
Any in Category	85.2%	90.6%	86.3%	80.8%	79.7%	85.5%	92.0%	93.5%	93.2%	93.8%
Tacrolimus	85.2%	90.6%	86.3%	80.8%	79.7%	85.5%	92.0%	93.0%	92.7%	93.8%
Everolimus	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.4%	0.5%	0.0%
Antimetabolites										
Any in Category	82.3%	75.5%	74.3%	85.6%	80.3%	83.8%	84.3%	94.3%	93.7%	94.9%
Mycophenolate Mofetil	80.5%	74.1%	74.0%	84.9%	78.9%	79.5%	74.3%	83.9%	79.1%	73.6%
Mycophenolate Sodium	0.0%	0.0%	0.0%	0.0%	1.1%	3.3%	9.6%	9.1%	13.1%	20.8%
Azathioprine	1.8%	1.4%	0.3%	0.6%	0.3%	1.0%	0.4%	1.3%	1.0%	0.6%
Leflunomide	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.5%	0.0%
mTOR Inhibitors										
Any in Category	8.3%	22.0%	19.4%	14.7%	16.3%	13.9%	15.7%	7.8%	7.9%	7.3%
Sirolimus	8.3%	22.0%	19.4%	14.7%	16.3%	13.9%	15.7%	7.8%	7.9%	7.3%
Everolimus	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

Individual drug percentages will not necessarily add up to the drug category percentages, as patients may be prescribed more than one drug within the same category.