

Table 8.6e
Immunosuppression Use for Maintenance
Prior to Discharge, 2000 to 2009
Recipients with Kidney-Pancreas Transplants

	Year of Transplant									
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Functioning Graft at Discharge	888	861	881	847	864	884	905	842	820	836
With Immunosuppression Info	886	861	881	844	851	879	891	833	817	821
Corticosteroids										
Any in Category	97.1%	93.3%	89.7%	83.8%	75.2%	71.0%	66.2%	65.4%	70.1%	69.9%
Prednisone	0.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Steroids	97.0%	93.3%	89.7%	83.8%	75.2%	71.0%	66.2%	65.4%	70.1%	69.9%
Cyclosporine										
Any in Category	18.8%	13.7%	9.8%	8.9%	6.2%	6.7%	6.4%	4.6%	4.4%	3.4%
Sandimmune	1.9%	0.9%	0.8%	0.1%	0.1%	0.2%	0.2%	0.2%	0.5%	0.2%
Neoral	16.1%	10.8%	6.8%	7.2%	5.3%	5.2%	5.4%	3.7%	3.7%	2.7%
Gengraf	0.8%	2.0%	1.9%	1.5%	0.8%	1.3%	0.8%	0.6%	0.2%	0.5%
Eon	0.0%	0.0%	0.2%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Tacrolimus										
Any in Category	82.3%	86.6%	90.2%	83.9%	84.6%	88.9%	92.4%	94.4%	94.0%	94.5%
Tacrolimus	82.3%	86.6%	90.2%	83.9%	84.6%	88.9%	92.3%	93.9%	93.3%	94.0%
Everolimus	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.5%	0.7%	0.5%
Antimetabolites										
Any in Category	84.9%	79.9%	82.2%	82.7%	83.7%	81.8%	85.1%	91.4%	91.1%	91.4%
Mycophenolate Mofetil	82.4%	78.9%	81.2%	81.6%	81.8%	74.6%	71.6%	73.5%	72.5%	64.7%
Mycophenolate Sodium	0.0%	0.0%	0.0%	0.0%	1.5%	6.9%	12.3%	17.2%	18.5%	25.9%
Azathioprine	2.5%	1.0%	1.0%	1.1%	0.4%	0.2%	1.1%	0.7%	0.1%	0.7%
Cytoxin	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
mTOR Inhibitors										
Any in Category	13.2%	21.4%	18.8%	19.3%	16.8%	15.4%	16.3%	10.3%	8.0%	6.1%
Sirolimus	13.0%	21.4%	18.8%	19.2%	16.8%	15.4%	16.3%	10.3%	8.0%	6.1%
Everolimus	0.2%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

Individual drug percentages will not necessarily add up to the drug category percentages, as patients may be prescribed more than one drug within the same category.