

**Table 8.6f**  
**Immunosuppression Use for Maintenance by Regimen**  
**One Year Following Transplantation, 1999 to 2008**  
**Recipients with Kidney-Pancreas Transplants**

	Transplant Year									
	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
<b>Transplants</b>	941	915	891	906	871	881	903	924	864	837
<b>Functioning Graft at 1 Year PostTx</b>	859	839	818	821	785	793	825	832	790	765
<b>With Maintenance Use Recorded at 1 Year PostTx</b>										
CyA	0.2%	0.0%	0.1%	0.3%	0.0%	0.0%	0.2%	0.1%	0.0%	0.0%
Tac	0.5%	0.1%	1.5%	2.6%	1.7%	2.8%	1.8%	2.0%	1.3%	1.3%
--- + MMF/MPA	0.0%	0.1%	0.1%	0.1%	0.4%	1.0%	0.2%	0.1%	0.0%	0.1%
CyA + MMF/MPA	0.0%	0.0%	0.0%	0.3%	0.5%	0.1%	1.0%	0.4%	0.9%	0.7%
Tac + MMF/MPA	1.0%	2.7%	2.1%	4.9%	6.3%	9.8%	14.2%	20.1%	20.4%	17.9%
--- + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%
CyA + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%
Tac + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.1%	0.1%	0.0%	0.1%
Tac + MMF/MPA + OtherAntimet	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%
--- + Siro/Evero	0.0%	0.1%	0.1%	0.4%	0.3%	0.0%	0.0%	0.0%	0.0%	0.3%
CyA + Siro/Evero	0.0%	0.2%	0.1%	0.0%	1.0%	2.2%	1.8%	2.3%	2.0%	2.1%
Tac + Siro/Evero	0.0%	3.0%	5.3%	7.3%	8.2%	6.0%	8.2%	3.3%	3.2%	2.6%
--- + MMF/MPA + Siro/Evero	0.0%	0.0%	0.1%	0.0%	1.5%	3.9%	1.5%	0.9%	0.7%	0.8%
CyA + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.3%	0.1%	0.0%	0.0%	0.0%
Tac + MMF/MPA + Siro/Evero	0.0%	0.1%	1.5%	0.9%	1.2%	1.0%	1.3%	3.3%	3.8%	1.9%
Tac + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.1%	0.0%	0.0%
--- + Steroids	0.2%	0.4%	0.1%	1.3%	0.5%	0.5%	0.1%	0.2%	0.3%	0.3%
CyA + Steroids	1.9%	1.7%	1.1%	0.5%	0.4%	0.1%	0.1%	0.2%	0.1%	0.3%
Tac + Steroids	6.0%	4.0%	5.3%	4.4%	5.0%	2.7%	4.5%	3.7%	4.6%	5.0%
--- + MMF/MPA + Steroids	0.4%	0.4%	0.4%	0.5%	0.4%	0.8%	0.6%	0.2%	0.4%	0.9%
CyA + MMF/MPA + Steroids	18.6%	11.5%	8.3%	8.8%	6.8%	3.2%	2.3%	2.0%	2.6%	1.9%
Tac + MMF/MPA + Steroids	57.6%	56.0%	51.9%	50.7%	52.1%	55.1%	53.9%	53.1%	53.8%	59.2%
--- + OtherAntimet + Steroids	0.0%	0.0%	0.0%	0.0%	0.1%	0.1%	0.0%	0.0%	0.0%	0.0%
CyA + OtherAntimet + Steroids	0.1%	0.4%	0.1%	0.4%	0.0%	0.3%	0.0%	0.0%	0.3%	0.0%
Tac + OtherAntimet + Steroids	3.7%	1.7%	0.8%	1.5%	1.7%	1.0%	0.6%	1.2%	0.9%	1.6%
--- + Siro/Evero + Steroids	0.2%	0.5%	0.4%	0.3%	0.6%	0.3%	0.0%	0.1%	0.3%	0.0%
CyA + Siro/Evero + Steroids	0.5%	3.0%	2.5%	1.0%	1.5%	0.5%	0.6%	0.6%	0.8%	0.1%
Tac + Siro/Evero + Steroids	1.5%	6.7%	12.8%	10.0%	5.5%	5.2%	4.2%	3.2%	2.5%	1.6%
Other Regimen	7.6%	7.3%	5.4%	4.0%	4.1%	3.2%	2.3%	2.3%	1.1%	1.2%

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

Percentages are calculated based on the total number of patients with maintenance use at one year following transplantation. They sum to 100 percent for each column.

---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.

MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.

"Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.