

Table 9.6d
Immunosuppression Use for Maintenance by Regimen
Prior to Discharge, 2000 to 2009
Recipients with Liver Transplants

	Transplant Year									
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Functioning Graft at Discharge	4,324	4,573	4,778	5,121	5,623	5,865	6,127	6,000	5,880	5,920
With Immunosuppression Info	4,310	4,567	4,775	5,116	5,595	5,796	6,062	5,965	5,840	5,909
CyA	0.2%	0.7%	0.2%	0.4%	0.2%	0.2%	0.1%	0.1%	0.2%	0.2%
Tac	1.5%	3.4%	4.8%	7.6%	9.8%	10.0%	5.6%	4.8%	3.7%	3.9%
--- + MMF/MPA	0.0%	0.1%	0.1%	0.3%	0.3%	0.2%	0.3%	0.2%	0.1%	0.2%
CyA + MMF/MPA	0.3%	0.2%	0.2%	0.4%	0.8%	0.6%	0.9%	1.2%	0.9%	0.8%
Tac + MMF/MPA	2.5%	2.6%	2.8%	6.5%	6.3%	8.5%	12.9%	15.6%	13.7%	12.4%
--- + OtherAntimet	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
CyA + OtherAntimet	0.0%	0.0%	0.1%	0.5%	0.5%	0.0%	0.0%	0.0%	0.0%	0.0%
Tac + OtherAntimet	0.1%	0.1%	0.0%	0.3%	0.3%	0.3%	0.1%	0.2%	0.0%	0.1%
CyA + MMF/MPA + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Tac + MMF/MPA + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
--- + Siro/Evero	0.0%	0.0%	0.1%	0.1%	0.1%	0.2%	0.1%	0.1%	0.0%	0.0%
CyA + Siro/Evero	0.6%	0.4%	0.0%	0.0%	0.0%	0.3%	0.6%	0.5%	0.3%	0.3%
Tac + Siro/Evero	1.0%	1.1%	0.4%	0.2%	0.4%	0.1%	0.2%	0.1%	0.1%	0.1%
--- + MMF/MPA + Siro/Evero	0.0%	0.0%	0.1%	0.2%	0.1%	0.1%	0.2%	0.2%	0.3%	0.2%
CyA + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%
Tac + MMF/MPA + Siro/Evero	0.0%	0.2%	0.1%	0.2%	0.1%	0.1%	0.1%	0.1%	0.0%	0.0%
--- + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
CyA + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Tac + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
--- + Steroids	2.8%	2.2%	0.8%	0.8%	1.4%	0.8%	0.6%	0.5%	1.1%	2.2%
CyA + Steroids	6.1%	2.6%	2.1%	1.4%	1.0%	1.1%	0.5%	0.7%	0.4%	0.6%
Tac + Steroids	32.1%	32.3%	35.4%	29.4%	26.2%	24.1%	21.5%	19.5%	16.1%	16.1%
--- + MMF/MPA + Steroids	0.6%	0.9%	0.9%	0.9%	1.2%	0.9%	1.0%	0.7%	1.0%	1.2%
CyA + MMF/MPA + Steroids	6.6%	5.5%	4.6%	3.8%	3.3%	3.1%	3.0%	3.0%	3.5%	3.4%
Tac + MMF/MPA + Steroids	30.5%	35.1%	36.6%	39.5%	41.4%	44.8%	48.1%	49.2%	55.3%	54.0%
--- + OtherAntimet + Steroids	0.0%	0.0%	0.0%	0.3%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%
CyA + OtherAntimet + Steroids	2.6%	1.2%	1.2%	0.1%	0.5%	0.2%	0.1%	0.1%	0.1%	0.1%
Tac + OtherAntimet + Steroids	1.9%	1.4%	2.0%	1.8%	0.6%	0.1%	0.1%	0.2%	0.2%	0.7%
--- + Siro/Evero + Steroids	0.4%	0.4%	0.4%	0.2%	0.3%	0.2%	0.2%	0.1%	0.1%	0.2%
CyA + Siro/Evero + Steroids	0.4%	0.3%	0.6%	0.5%	0.4%	0.3%	0.1%	0.1%	0.2%	0.1%
Tac + Siro/Evero + Steroids	4.5%	4.6%	2.6%	1.0%	1.4%	0.9%	1.0%	0.4%	0.3%	0.2%
Other Regimen	5.0%	4.7%	3.9%	3.4%	3.1%	2.8%	2.4%	2.2%	2.3%	2.8%

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

Percentages in each column sum to 100 percent.

---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.

MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.

"Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.