

Table 9.6g
Immunosuppression Use for Maintenance Between Discharge
and One Year Following Transplantation, 1999 to 2008
Recipients with Liver Transplants

	Year of Transplant									
	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Transplants	4,755	5,001	5,195	5,332	5,673	6,171	6,444	6,651	6,494	6,319
Tx with Follow-up Immunosuppression Info	3,643	3,890	4,090	4,304	4,566	4,940	5,054	5,303	5,155	4,973
Corticosteroids										
Any in Category	89.3%	90.5%	89.5%	87.2%	78.5%	75.4%	75.3%	72.0%	68.2%	69.3%
Steroids	89.3%	90.5%	89.5%	87.2%	78.5%	75.4%	75.3%	72.0%	68.2%	69.3%
Cyclosporine										
Any in Category	30.4%	23.4%	18.4%	14.0%	11.8%	11.6%	10.4%	9.9%	10.0%	10.1%
Cyclosporin	0.1%	0.1%	0.0%	0.0%	0.0%	0.0%	0.1%	0.1%	0.1%	0.3%
Sandimmune	7.0%	5.4%	3.7%	2.0%	0.7%	0.9%	1.1%	1.2%	1.3%	1.1%
Neoral	23.3%	17.3%	13.3%	9.8%	8.9%	9.2%	7.8%	6.7%	7.1%	6.5%
Gengraf	0.1%	0.5%	1.3%	2.0%	1.8%	1.3%	1.3%	1.7%	1.2%	1.7%
Eon	0.0%	0.1%	0.2%	0.2%	0.3%	0.2%	0.1%	0.2%	0.2%	0.5%
Tacrolimus										
Any in Category	76.6%	84.1%	89.1%	90.5%	90.5%	89.7%	91.7%	91.0%	90.7%	89.6%
Tacrolimus	76.6%	84.1%	89.1%	90.5%	90.5%	89.7%	91.7%	90.8%	90.5%	89.3%
Everolimus	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.2%	0.3%	0.3%
Antimetabolites										
Any in Category	49.3%	52.2%	53.5%	53.7%	55.1%	59.0%	66.3%	69.8%	69.9%	72.9%
Mycophenolate Mofetil	40.8%	46.2%	48.8%	49.3%	52.4%	55.5%	62.2%	64.4%	63.0%	65.5%
Mycophenolate Sodium	0.0%	0.0%	0.0%	0.0%	0.0%	0.6%	2.8%	4.4%	5.4%	6.2%
Azathioprine	8.3%	5.9%	4.6%	4.3%	2.7%	2.9%	1.2%	1.0%	1.4%	1.2%
Leflunomide	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Cytosan	0.1%	0.1%	0.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
mTOR Inhibitors										
Any in Category	3.2%	15.1%	16.6%	13.8%	12.2%	14.1%	13.8%	11.9%	9.9%	9.0%
Sirolimus	3.1%	15.1%	16.6%	13.8%	12.2%	14.1%	13.8%	11.8%	9.9%	8.8%
Everolimus	0.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.2%

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

Percentages are calculated based on the total number of patients with any immunosuppression use between discharge and one year following transplantation. Individual drug percentages will not necessarily add up to the drug category percentages, as patients may be prescribed more than one drug within the same category.