Table 7.6b
Immunosuppression Use for Induction by Discharge Regimen, 2001 to 2005
Recipients with Pancreas After Kidney (PAK) Transplants

	Discharge Regimen (w/ or w/o Steroids Use)										
	CyA	Tac	CyA+ MMF	Tac+ MMF	CyA+ Aza	Tac+ Aza	Siro+ MMF	CyA+ Siro	Tac+ Siro	Other	All
Functioning Graft with Immunosuppression Info at Discharge	6	74	93	1,046	1	8	22	27	172	159	1,608
% of All Regimens	0.4%	4.6%	5.8%	65.0%	0.1%	0.5%	1.4%	1.7%	10.7%	9.9%	100.0%
Induction Drug											
Atgam/NRATG/NRATS	0.0%	0.0%	0.0%	2.3%	0.0%	0.0%	4.5%	0.0%	1.2%	1.3%	1.8%
OKT3	0.0%	0.0%	0.0%	0.5%	0.0%	0.0%	0.0%	0.0%	1.2%	2.5%	0.7%
Thymoglobulin	50.0%	13.5%	55.9%	55.3%	0.0%	50.0%	50.0%	85.2%	61.0%	91.8%	58.0%
Zenapax	0.0%	2.7%	12.9%	17.2%	0.0%	12.5%	0.0%	0.0%	26.7%	22.6%	17.2%
Simulect	0.0%	1.4%	16.1%	7.6%	0.0%	0.0%	4.5%	14.8%	6.4%	11.3%	8.0%
Campath	16.7%	18.9%	8.6%	5.3%	0.0%	25.0%	22.7%	0.0%	9.3%	50.9%	11.3%
No Induction Drugs Recorded	50.0%	63.5%	20.4%	21.7%	100.0%	25.0%	36.4%	3.7%	19.8%	14.5%	22.7%

Source: OPTN/SRTR Data as of May 1, 2006.
Notes:
Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.
CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.