Table 10.6b
Immunosuppression Use for Induction by Discharge Regimen, 2001 to 2005
Recipients with Intestine Transplants

	Discharge Regimen (w/ or w/o Steroids Use)						
	CyA	Tac	Tac+ MMF	Tac+ Aza	Tac+ Siro	Other	All
Functioning Graft with Immunosuppression Info at Discharge	1	413	45	6	65	26	556
% of All Regimens	0.2%	74.3%	8.1%	1.1%	11.7%	4.7%	100.0%
Induction Drug							
Atgam/NRATG/NRATS	0.0%	0.2%	8.9%	0.0%	1.5%	15.4%	1.8%
OKT3	0.0%	2.9%	0.0%	0.0%	0.0%	26.9%	3.4%
Thymoglobulin	0.0%	24.5%	20.0%	0.0%	26.2%	100.0%	27.5%
Zenapax	0.0%	8.0%	53.3%	100.0%	15.4%	73.1%	16.5%
Simulect	0.0%	2.4%	0.0%	0.0%	30.8%	26.9%	6.7%
Campath	0.0%	15.3%	6.7%	0.0%	0.0%	34.6%	13.5%
No Induction Drugs Recorded	100.0%	48.2%	17.8%	0.0%	26.2%	38.5%	42.3%

Source: OPTN/SRTR Data as of May 1, 2006.
Notes:
Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.
CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.