Table 11.6b
Immunosuppression Use for Induction by Discharge Regimen, 2001 to 2005
Recipients with Heart Transplants

	Discharge Regimen (w/ or w/o Steroids Use)										All
	CyA	Tac	CyA+ MMF	Tac+ MMF	CyA+ Aza	Tac+ Aza	Siro+ MMF	CyA+ Siro	Tac+ Siro	Other	
Functioning Graft with Immunosuppression Info at Discharge	151	302	4,066	3,094	570	123	69	65	229	908	9,577
% of All Regimens	1.6%	3.2%	42.5%	32.3%	6.0%	1.3%	0.7%	0.7%	2.4%	9.5%	100.0%
Induction Drug											
Atgam/NRATG/NRATS	9.9%	6.3%	7.6%	5.4%	6.7%	12.2%	4.3%	4.6%	8.7%	18.5%	7.9%
OKT3	4.6%	2.3%	5.7%	2.5%	14.9%	5.7%	1.4%	0.0%	4.8%	11.0%	5.5%
Thymoglobulin	5.3%	6.3%	12.6%	15.3%	16.7%	21.1%	20.3%	4.6%	20.1%	27.3%	15.1%
Zenapax	4.0%	1.3%	17.2%	11.1%	3.2%	3.3%	14.5%	75.4%	0.4%	34.1%	15.1%
Simulect	7.9%	7.3%	10.7%	9.6%	3.2%	1.6%	7.2%	4.6%	5.7%	13.3%	9.7%
Campath	0.0%	1.0%	0.0%	1.3%	0.0%	0.0%	0.0%	0.0%	0.0%	0.8%	0.5%
No Induction Drugs Recorded	70.2%	77.8%	48.0%	59.0%	57.7%	56.9%	52.2%	12.3%	62.9%	42.5%	53.2%

Source: OPTN/SRTR Data as of May 1, 2006.
Notes:
Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.
CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.