Table 2.15
Deceased Donor Organ Utilization, 1996 to 2005, Liver

	Year									
	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total # Donors of Any Organ	5,418	5,479	5,793	5,824	5,985	6,080	6,190	6,457	7,150	7,593
# Livers Recovered	4,463	4,600	4,843	4,947	4,997	5,106	5,294	5,682	6,320	6,690
ORPD (1)	0.82	0.84	0.84	0.85	0.83	0.84	0.86	0.88	0.88	0.88
# Livers Transplanted	3,976	4,037	4,349	4,425	4,505	4,597	4,893	5,263	5,778	6,042
OTPD (2)	0.73	0.74	0.75	0.76	0.75	0.76	0.79	0.82	0.81	0.80
% Transplated	89.1%	87.8%	89.8%	89.4%	90.2%	90.0%	92.4%	92.6%	91.4%	90.3%
Donor with Brain Death(DBD)										
Total # Donors of Any Organ	5,362	5,410	5,726	5,748	5,880	5,935	6,033	6,237	6,829	7,134
# Livers Recovered	4,447	4,574	4,809	4,915	4,950	5,028	5,205	5,547	6,113	6,361
ORPD (1)	0.83	0.85	0.84	0.86	0.84	0.85	0.86	0.89	0.90	0.89
# Livers Transplanted	3,964	4,021	4,326	4,404	4,472	4,537	4,824	5,163	5,613	5,805
OTPD (2)	0.74	0.74	0.76	0.77	0.76	0.76	0.80	0.83	0.82	0.81
% Transplated	89.1%	87.9%	90.0%	89.6%	90.3%	90.2%	92.7%	93.1%	91.8%	91.3%
Donors After Cardiac Death(DCD)										
Total # Donors of Any Organ	56	69	67	76	105	145	157	220	321	459
# Livers Recovered	16	26	34	32	47	78	89	135	207	329
ORPD (1)	0.29	0.38	0.51	0.42	0.45	0.54	0.57	0.61	0.64	0.72
# Livers Transplanted	12	16	23	21	33	60	69	100	165	237
OTPD (2)	0.21	0.23	0.34	0.28	0.31	0.41	0.44	0.45	0.51	0.52
% Transplated	75.0%	61.5%	67.6%	65.6%	70.2%	76.9%	77.5%	74.1%	79.7%	72.0%
Liver Donors	4,463	4,600	4,843	4,947	4,997	5,106	5,294	5,682	6,320	6,690
Liver Donors (%)	82.4%	84.0%	83.6%	84.9%	83.5%	84.0%	85.5%	88.0%	88.4%	88.1%
Liver Donors, Other Organs Recovered										
Liver Only	205	216	273	275	320	378	382	532	655	699
Liver & Kidney or Pancreas Only	1,923	2,041	2,214	2,404	2,425	2,412	2,606	2,881	3,302	3,419
Liver & Intestine Only	1	5	6	4	4	3	3	5	7	5
Liver & Thoracic Only	83	75	76	79	73	69	71	84	81	100
Liver & Other Combinations	2,251	2,263	2,274	2,185	2,175	2,244	2,232	2,180	2,275	2,467
Liver Donors, Other Organs Recovered (%)										
Liver Only	4.6%	4.7%	5.6%	5.6%	6.4%	7.4%	7.2%	9.4%	10.4%	10.4%

(Continued)

Source: OPTN/SRTR Data as of May 1, 2006.
Notes:
(-) = None in category.
(1) ORPD: Organs Recovered Per Donor of any organ; (2) OTPD: Organs Transplanted Per Donor of any organ. Organs divided into segments (liver, lung, pancreas, intestine) are counted only once for that organ.
Includes organs recovered for transplant but not transplanted, as well as organs transplanted.
Donors are divided into two mutually exclusive and complete categories. DCDs include patients whose hearts have irreversibly stopped beating, and may also have been known as Non-Heart Beating Donors. DBDs have suffered brain death.

Table 2.15 (Continued)
Deceased Donor Organ Utilization, 1996 to 2005, Liver

	Year									
	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Liver Donors, Other Organs Recovered (%)										
Liver & Kidney or Pancreas Only	43.1%	44.4%	45.7%	48.6%	48.5%	47.2%	49.2%	50.7%	52.2%	51.1%
Liver & Intestine Only	0.0%	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%
Liver & Thoracic Only	1.9%	1.6%	1.6%	1.6%	1.5%	1.4%	1.3%	1.5%	1.3%	1.5%
Liver & Other Combinations	50.4%	49.2%	47.0%	44.2%	43.5%	43.9%	42.2%	38.4%	36.0%	36.9%

Source: OPTN/SRTR Data as of May 1, 2006.
Notes:
(-) = None in category.
(1) ORPD: Organs Recovered Per Donor of any organ; (2) OTPD: Organs Transplanted Per Donor of any organ. Organs divided into segments (liver, lung, pancreas, intestine) are counted only once for that organ.
Includes organs recovered for transplant but not transplanted, as well as organs transplanted.
Donors are divided into two mutually exclusive and complete categories. DCDs include patients whose hearts have irreversibly stopped beating, and may also have been known as Non-Heart Beating Donors. DBDs have suffered brain death.