Table 6.6b
Immunosuppression Use for Induction by Discharge Regimen, 2001 to 2005
Recipients with Pancreas Transplant Alone (PTA) Transplants

	Discharge Regimen (w/ or w/o Steroids Use)								All
	Tac	CyA+ MMF	Tac+ MMF	Tac+ Aza	Siro+ MMF	CyA+ Siro	Tac+ Siro	Other	
Functioning Graft with Immunosuppression Info at Discharge	227	13	377	5	4	3	72	99	800
% of All Regimens	28.4%	1.6%	47.1%	0.6%	0.5%	0.4%	9.0%	12.4%	100.0%
Induction Drug									
Atgam/NRATG/NRATS	0.0%	0.0%	1.6%	0.0%	0.0%	0.0%	0.0%	0.0%	0.8%
OKT3	3.1%	0.0%	1.1%	0.0%	0.0%	0.0%	0.0%	3.0%	1.8%
Thymoglobulin	26.0%	61.5%	66.6%	60.0%	75.0%	66.7%	43.1%	81.8%	54.8%
Zenapax	9.7%	30.8%	24.1%	60.0%	0.0%	0.0%	16.7%	17.2%	18.6%
Simulect	0.4%	23.1%	3.7%	0.0%	0.0%	0.0%	2.8%	6.1%	3.3%
Campath	18.9%	15.4%	9.0%	20.0%	25.0%	0.0%	38.9%	80.8%	23.6%
No Induction Drugs Recorded	44.5%	15.4%	17.2%	0.0%	25.0%	33.3%	19.4%	11.1%	24.4%

Source: OPTN/SRTR Data as of May 1, 2006.
Notes:
Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.
CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.