Table 8.6f
Immunosuppression Use for Maintenance by Regimen
One Year Following Transplantation, 1995 to 2004
Recipients with Kidney-Pancreas Transplants

	Transplant Year									
	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
Transplants	919	861	853	972	942	915	891	905	871	881
Functioning Graft at 1 Year PostTx	825	774	781	871	860	841	818	822	783	784
With Maintenance Use Recorded at 1 Year PostTx	79	83	179	631	840	810	798	797	777	769
--- + MMF/MPA	7.6%	0.0%	0.0%	0.0%	0.0%	0.1%	0.1%	0.1%	0.4%	1.0%
--- + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	1.5%	4.0%
--- + MMF/MPA + Steroids	0.0%	0.0%	0.6%	0.6%	0.4%	0.4%	0.4%	0.5%	0.4%	0.8%
--- + OtherAntimet + Steroids	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.1%
--- + Siro/Evero	0.0%	0.0%	0.0%	0.2%	0.0%	0.1%	0.1%	0.4%	0.3%	0.0%
--- + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.0%	0.2%	0.5%	0.3%	0.3%	0.6%	0.3%
--- + Steroids	0.0%	0.0%	0.0%	0.3%	0.2%	0.4%	0.1%	1.3%	0.6%	0.5%
CyA	5.1%	0.0%	0.0%	0.2%	0.2%	0.0%	0.1%	0.3%	0.0%	0.0%
CyA + MMF/MPA	1.3%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.3%	0.5%	0.1%
CyA + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.3%
CyA + MMF/MPA + Steroids	12.7%	20.5%	29.1%	24.1%	18.6%	11.5%	8.3%	8.8%	6.8%	3.3%
CyA + OtherAntimet + Steroids	22.8%	9.6%	5.0%	1.9%	0.1%	0.4%	0.1%	0.4%	0.0%	0.3%
CyA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.2%	0.1%	0.0%	1.0%	2.2%
CyA + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.0%	0.5%	3.0%	2.5%	1.1%	1.5%	0.5%
CyA + Steroids	3.8%	6.0%	0.6%	1.6%	1.9%	1.9%	1.1%	0.5%	0.4%	0.1%
Tac	0.0%	1.2%	1.1%	1.0%	0.5%	0.1%	1.5%	2.9%	1.7%	2.9%
Tac + MMF/MPA	2.5%	0.0%	0.0%	0.6%	1.0%	2.7%	2.1%	4.9%	6.4%	9.8%
Tac + MMF/MPA + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%
Tac + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	1.5%	0.9%	1.0%	1.0%
Tac + MMF/MPA + Steroids	11.4%	39.8%	48.0%	54.8%	57.6%	56.0%	52.0%	50.4%	52.0%	54.5%
Tac + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%
Tac + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%
Tac + OtherAntimet + Steroids	5.1%	7.2%	2.2%	3.6%	3.6%	1.7%	0.8%	1.5%	1.5%	1.0%
Tac + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	3.0%	5.3%	7.3%	8.2%	6.1%
Tac + Siro/Evero + Steroids	0.0%	0.0%	0.6%	0.0%	1.5%	6.5%	12.9%	9.9%	5.5%	5.2%
Tac + Steroids	5.1%	6.0%	5.6%	5.1%	6.0%	4.0%	5.1%	4.3%	5.0%	2.6%
Other Regimen	22.8%	9.6%	7.3%	6.0%	7.7%	7.3%	5.4%	4.1%	4.1%	3.3%

Source: OPTN/SRTR Data as of May 1, 2006.

Notes:

Percentages are calculated based on the total number of patients with maintenance use at one year following transplantation. They sum to 100 percent for each column.

---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.

MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.

"Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.