Table 10.6f
Immunosuppression Use for Maintenance by Regimen
One Year Following Transplantation, 1995 to 2004
Recipients with Intestine Transplants

	Transplant Year									
	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
Transplants	46	45	68	70	73	82	112	107	116	152
Functioning Graft at 1 Year PostTx	22	27	31	34	37	55	69	73	88	113
With Maintenance Use Recorded at 1 Year PostTx	6	2	7	21	36	55	69	72	87	112
--- + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	1.4%	1.4%	1.1%	0.9%
--- + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.0%	0.0%	1.8%	0.0%	0.0%	1.1%	3.6%
--- + Steroids	0.0%	0.0%	0.0%	0.0%	0.0%	1.8%	0.0%	1.4%	3.4%	1.8%
CyA	16.7%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
CyA + Steroids	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.9%
Tac	33.3%	0.0%	57.1%	14.3%	2.8%	3.6%	26.1%	20.8%	39.1%	28.6%
Tac + MMF/MPA	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	1.4%	1.1%	0.0%
Tac + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.9%
Tac + MMF/MPA + Steroids	16.7%	0.0%	14.3%	23.8%	11.1%	9.1%	2.9%	4.2%	6.9%	6.3%
Tac + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	2.7%
Tac + OtherAntimet + Steroids	0.0%	50.0%	0.0%	4.8%	11.1%	0.0%	2.9%	0.0%	0.0%	1.8%
Tac + Siro/Evero	0.0%	0.0%	0.0%	0.0%	2.8%	0.0%	1.4%	1.4%	1.1%	3.6%
Tac + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.0%	8.3%	21.8%	7.2%	20.8%	4.6%	8.9%
Tac + Steroids	16.7%	50.0%	28.6%	57.1%	61.1%	61.8%	58.0%	45.8%	41.4%	40.2%
Other Regimen	16.7%	0.0%	0.0%	0.0%	2.8%	0.0%	0.0%	2.8%	0.0%	0.0%

Source: OPTN/SRTR Data as of May 1, 2006.

Notes:

Percentages are calculated based on the total number of patients with maintenance use at one year following transplantation. They sum to 100 percent for each column.

---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.

MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.

"Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.