Table 5.6b
Immunosuppression Use for Induction by Discharge Regimen, 2001 to 2005
Recipients with Kidney Transplants

	Discharge Regimen (w/ or w/o Steroids Use)										All
	CyA	Tac	CyA+ MMF	Tac+ MMF	CyA+ Aza	Tac+ Aza	Siro+ MMF	CyA+ Siro	Tac+ Siro	Other	
Functioning Graft with Immunosuppression Info at Discharge	1,007	3,076	14,412	40,453	529	604	2,043	2,095	4,178	5,190	73,587
% of All Regimens	1.4%	4.2%	19.6%	55.0%	0.7%	0.8%	2.8%	2.8%	5.7%	7.1%	100.0%
Induction Drug											
Atgam/NRATG/NRATS	1.5%	0.7%	1.6%	1.9%	6.6%	6.1%	2.1%	1.4%	1.1%	2.4%	1.8%
OKT3	0.3%	0.4%	0.4%	0.6%	0.6%	0.3%	0.2%	0.2%	0.6%	1.0%	0.6%
Thymoglobulin	14.2%	24.3%	19.8%	34.3%	28.2%	30.1%	32.1%	39.3%	36.4%	40.8%	31.3%
Zenapax	8.2%	4.6%	17.7%	12.8%	5.7%	10.3%	14.7%	2.7%	7.4%	12.4%	12.7%
Simulect	15.4%	15.4%	29.1%	18.5%	35.7%	28.3%	31.3%	21.6%	21.6%	30.5%	22.1%
Campath	0.4%	12.7%	2.3%	4.0%	0.0%	0.3%	2.1%	0.0%	0.6%	11.1%	4.1%
No Induction Drugs Recorded	61.3%	43.8%	31.0%	31.6%	25.0%	25.2%	21.7%	36.0%	34.8%	29.6%	32.2%

Source: OPTN/SRTR Data as of May 1, 2006.
Notes:
Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.
CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.