Table 1.9c
Immunosuppression Use by Organ
Maintenance Regimen at One Year Following Transplantation
2004 Transplants

	Organ								
	Kidney	PTA	PAK	KP	Liver	Intest-ine	Heart	Lung	Heart-Lung
Transplants	16,004	183	420	881	6,167	152	2,016	1,185	39
With Maintenance Use Recorded at 1 Year PostTx	13,958	138	317	769	4,861	112	1,709	973	28
Maintenance Regimen									
--- + MMF/MPA	0.2%	3.6%	3.2%	1.0%	0.3%	0.0%	0.2%	0.0%	0.0%
--- + MMF/MPA + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%
--- + MMF/MPA + Siro/Evero	1.3%	0.7%	0.3%	4.0%	1.3%	0.0%	0.2%	0.0%	0.0%
--- + MMF/MPA + Steroids	0.8%	0.0%	0.3%	0.8%	0.4%	0.0%	0.6%	0.2%	0.0%
--- + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.1%	0.0%	0.0%
--- + OtherAntimet + Steroids	0.0%	0.7%	0.0%	0.1%	0.0%	0.0%	0.0%	0.1%	0.0%
--- + Siro/Evero	0.2%	2.2%	0.3%	0.0%	2.0%	0.9%	0.0%	0.0%	0.0%
--- + Siro/Evero + Steroids	0.7%	1.4%	0.0%	0.3%	1.6%	3.6%	0.1%	0.2%	0.0%
--- + Steroids	0.3%	0.7%	0.0%	0.5%	0.5%	1.8%	0.4%	0.4%	0.0%
CyA	0.1%	0.0%	0.0%	0.0%	1.9%	0.0%	0.3%	0.1%	0.0%
CyA + MMF/MPA	2.0%	0.7%	0.9%	0.1%	2.3%	0.0%	8.5%	0.0%	0.0%
CyA + MMF/MPA + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%
CyA + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.3%	0.1%	0.0%	0.1%	0.0%	0.0%
CyA + MMF/MPA + Steroids	12.6%	1.4%	3.8%	3.3%	1.9%	0.0%	23.5%	7.6%	14.3%
CyA + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.2%	0.0%	1.4%	0.0%	0.0%
CyA + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%
CyA + OtherAntimet + Steroids	0.4%	0.0%	0.0%	0.3%	0.3%	0.0%	2.7%	6.2%	10.7%
CyA + Siro/Evero	0.9%	0.0%	0.6%	2.2%	0.3%	0.0%	0.5%	0.0%	0.0%
CyA + Siro/Evero + Steroids	1.5%	0.7%	0.6%	0.5%	0.1%	0.0%	1.1%	0.3%	0.0%
CyA + Steroids	1.2%	0.0%	0.0%	0.1%	1.7%	0.9%	0.9%	1.2%	0.0%
Tac	2.2%	15.2%	5.4%	2.9%	28.3%	28.6%	1.2%	0.1%	0.0%
Tac + MMF/MPA	14.1%	12.3%	8.5%	9.8%	15.3%	0.0%	10.8%	0.4%	0.0%
Tac + MMF/MPA + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Tac + MMF/MPA + Siro/Evero	0.7%	2.9%	1.3%	1.0%	0.2%	0.9%	0.3%	0.0%	0.0%
Tac + MMF/MPA + Steroids	42.9%	22.5%	50.5%	54.5%	15.4%	6.3%	32.1%	40.5%	35.7%
Tac + OtherAntimet	0.2%	0.7%	0.0%	0.1%	0.6%	2.7%	1.3%	0.2%	0.0%
Tac + OtherAntimet + Siro/Evero	0.0%	0.7%	0.0%	0.0%	0.0%	0.0%	0.1%	0.1%	0.0%
Tac + OtherAntimet + Steroids	1.0%	2.2%	0.9%	1.0%	0.5%	1.8%	1.5%	23.1%	21.4%
Tac + Siro/Evero	2.1%	8.0%	7.3%	6.1%	2.9%	3.6%	0.9%	0.0%	0.0%
Tac + Siro/Evero + Steroids	3.5%	2.9%	7.3%	5.2%	1.5%	8.9%	2.1%	4.2%	7.1%
Tac + Steroids	3.9%	12.3%	2.2%	2.6%	18.1%	40.2%	2.6%	10.0%	10.7%
Other Regimen	7.2%	8.0%	6.6%	3.3%	1.8%	0.0%	6.3%	5.0%	0.0%

Source: OPTN/SRTR Data as of May 1, 2006.

Notes:

Percentages are calculated based on the total number of patients with functioning graft and having any immunosuppression information reported at 1yr post-transplant. They sum to 100 percent.

---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.

MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF/MPA.

"Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.