Table 12.6f
Immunosuppression Use for Maintenance by Regimen
One Year Following Transplantation, 1995 to 2004
Recipients with Lung Transplants

	Transplant Year									
	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
Transplants	893	836	948	893	922	977	1,083	1,054	1,099	1,185
Functioning Graft at 1 Year PostTx	634	564	675	630	667	735	805	832	895	982
With Maintenance Use Recorded at 1 Year PostTx	30	53	88	324	620	695	767	808	864	973
--- + MMF/MPA	0.0%	0.0%	0.0%	0.3%	0.3%	0.1%	0.0%	0.0%	0.0%	0.0%
--- + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.1%	0.0%	0.0%
--- + MMF/MPA + Steroids	0.0%	0.0%	0.0%	0.3%	0.0%	0.4%	0.3%	0.1%	0.2%	0.2%
--- + OtherAntimet	0.0%	1.9%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%
--- + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.2%	0.0%	0.0%
--- + OtherAntimet + Steroids	0.0%	0.0%	1.1%	0.3%	0.2%	0.4%	0.3%	0.2%	0.3%	0.1%
--- + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.3%	0.0%	0.3%	0.1%	0.1%	0.0%
--- + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.7%	0.2%	0.2%
--- + Steroids	0.0%	1.9%	3.4%	0.9%	0.5%	0.4%	0.5%	0.4%	0.7%	0.4%
CyA	0.0%	0.0%	0.0%	0.0%	0.0%	0.4%	0.0%	0.1%	0.0%	0.1%
CyA + MMF/MPA	0.0%	1.9%	1.1%	0.0%	0.0%	0.1%	0.4%	0.1%	0.3%	0.0%
CyA + MMF/MPA + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%
CyA + MMF/MPA + Steroids	6.7%	7.5%	17.0%	15.4%	17.4%	14.0%	14.0%	10.8%	8.9%	7.6%
CyA + OtherAntimet	3.3%	0.0%	0.0%	0.3%	0.2%	0.1%	0.1%	0.4%	0.2%	0.0%
CyA + OtherAntimet + Steroids	60.0%	30.2%	30.7%	32.4%	26.8%	22.9%	12.3%	8.2%	8.9%	6.2%
CyA + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.3%	1.8%	1.3%	0.9%	0.9%	0.5%	0.3%
CyA + Steroids	3.3%	9.4%	6.8%	4.3%	5.6%	3.5%	2.1%	1.6%	2.2%	1.2%
Tac	0.0%	0.0%	0.0%	0.3%	0.2%	0.0%	0.3%	0.6%	0.3%	0.1%
Tac + MMF/MPA	0.0%	1.9%	0.0%	0.6%	0.3%	1.3%	0.5%	1.0%	0.5%	0.4%
Tac + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%
Tac + MMF/MPA + Steroids	0.0%	13.2%	18.2%	17.6%	17.1%	26.6%	29.7%	32.7%	34.3%	40.5%
Tac + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.2%	0.1%	0.1%	1.2%	0.5%	0.2%
Tac + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.1%
Tac + OtherAntimet + Steroids	13.3%	18.9%	10.2%	17.9%	17.4%	16.3%	17.3%	20.0%	22.9%	23.1%
Tac + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.2%	0.1%	0.1%	0.2%	0.1%	0.0%
Tac + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.0%	3.2%	2.7%	3.5%	6.3%	3.9%	4.2%
Tac + Steroids	6.7%	3.8%	2.3%	5.6%	6.5%	5.5%	10.0%	7.3%	8.0%	10.0%
Other Regimen	6.7%	9.4%	9.1%	3.4%	1.9%	3.5%	7.0%	6.6%	6.5%	5.0%

Source: OPTN/SRTR Data as of May 1, 2006.

Notes:

Percentages are calculated based on the total number of patients with maintenance use at one year following transplantation. They sum to 100 percent for each column.

---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.

MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.

"Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.