Table 10.6g
Immunosuppression Use for Maintenance Between Discharge and One Year Following Transplantation, 1995 to 2004
Recipients with Intestine Transplants

	Year of Transplant									
	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
Transplants	46	45	68	70	73	82	112	107	116	152
Tx with Follow-up Immunosuppression Info	22	27	31	34	37	55	69	73	88	113
Corticosteroids										
Any in Category	90.9%	100.0%	93.5%	97.1%	100.0%	100.0%	85.5%	91.8%	83.0%	78.8%
Steroids	90.9%	100.0%	93.5%	97.1%	100.0%	100.0%	85.5%	91.8%	83.0%	78.8%
Cyclosporine										
Any in Category	13.6%	11.1%	6.5%	2.9%	2.7%	0.0%	0.0%	1.4%	0.0%	0.9%
Cyclosporin	4.5%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Sandimmmune	0.0%	0.0%	0.0%	0.0%	2.7%	0.0%	0.0%	1.4%	0.0%	0.0%
Neoral	9.1%	11.1%	6.5%	2.9%	0.0%	0.0%	0.0%	0.0%	0.0%	0.9%
Tacrolimus										
Any in Category	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	98.9%	97.3%
Tacrolimus	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	98.9%	97.3%
Antimetabolites										
Any in Category	54.5%	66.7%	29.0%	41.2%	37.8%	14.5%	15.9%	11.0%	10.2%	12.4%
Mycophenolate Mofetil	18.2%	44.4%	12.9%	35.3%	21.6%	14.5%	2.9%	9.6%	10.2%	8.0%
Azathioprine	36.4%	22.2%	16.1%	5.9%	16.2%	0.0%	13.0%	1.4%	0.0%	4.4%
mTOR Inhibitors										
Any in Category	0.0%	0.0%	0.0%	0.0%	21.6%	69.1%	24.6%	28.8%	15.9%	22.1%
Sirolimus	0.0%	0.0%	0.0%	0.0%	21.6%	69.1%	24.6%	28.8%	15.9%	22.1%
Everolimus	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%

Source: OPTN/SRTR Data as of May 1, 2006.
Notes:
Percentages are calculated based on the total number of patients with any immunosuppression use between discharge and one year following transplantation.
Individual drug percentages will not necessarily add up to the drug category percentages, as patients may be prescribed more than one drug within the same category.