Table 12.6b
Immunosuppression Use for Induction by Discharge Regimen, 2001 to 2005
Recipients with Lung Transplants

	Discharge Regimen (w/ or w/o Steroids Use)										All
	CyA	Tac	CyA+ MMF	Tac+ MMF	CyA+ Aza	Tac+ Aza	Siro+ MMF	CyA+ Siro	Tac+ Siro	Other	
Functioning Graft with Immunosuppression Info at Discharge	76	285	686	1,655	957	1,282	10	5	32	272	5,260
% of All Regimens	1.4%	5.4%	13.0%	31.5%	18.2%	24.4%	0.2%	0.1%	0.6%	5.2%	100.0%
Induction Drug											
Atgam/NRATG/NRATS	0.0%	2.8%	2.0%	7.9%	8.3%	4.5%	0.0%	0.0%	3.1%	14.0%	6.2%
OKT3	1.3%	0.0%	1.5%	0.0%	3.1%	0.0%	0.0%	0.0%	0.0%	2.2%	0.9%
Thymoglobulin	5.3%	1.8%	3.6%	5.9%	5.7%	2.5%	0.0%	0.0%	0.0%	15.4%	5.0%
Zenapax	3.9%	2.1%	17.8%	14.3%	8.9%	17.7%	20.0%	0.0%	6.3%	30.5%	14.6%
Simulect	5.3%	4.2%	13.7%	19.3%	10.6%	27.3%	30.0%	40.0%	15.6%	47.4%	19.4%
Campath	1.3%	15.1%	0.0%	4.0%	0.0%	0.0%	0.0%	0.0%	3.1%	4.4%	2.3%
No Induction Drugs Recorded	82.9%	74.4%	61.8%	49.1%	64.3%	48.7%	50.0%	60.0%	71.9%	58.8%	55.9%

Source: OPTN/SRTR Data as of May 1, 2006.
Notes:
Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.
CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.