Table 5.6f
Immunosuppression Use for Maintenance by Regimen
One Year Following Transplantation, 1995 to 2004
Recipients with Kidney Transplants

	Transplant Year									
	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
Transplants	11,081	11,397	11,701	12,451	12,758	13,611	14,265	14,778	15,136	16,004
Functioning Graft at 1 Year PostTx	9,463	9,905	10,233	10,825	11,043	11,831	12,723	13,254	13,572	14,302
With Maintenance Use Recorded at 1 Year PostTx	636	808	1,537	6,744	10,468	11,216	12,493	12,935	13,222	13,958
--- + MMF/MPA	5.7%	1.5%	1.7%	0.3%	0.1%	0.0%	0.1%	0.1%	0.1%	0.2%
--- + MMF/MPA + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
--- + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.3%	0.5%	1.1%	1.3%
--- + MMF/MPA + Steroids	0.6%	0.9%	0.8%	1.2%	1.5%	0.9%	0.9%	0.9%	0.8%	0.8%
--- + OtherAntimet	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
--- + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
--- + OtherAntimet + Steroids	0.5%	0.6%	0.1%	0.2%	0.2%	0.1%	0.1%	0.1%	0.1%	0.0%
--- + Siro/Evero	0.8%	0.0%	0.3%	0.1%	0.1%	0.2%	0.2%	0.1%	0.1%	0.2%
--- + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.2%	0.3%	0.5%	0.7%	0.7%	0.6%	0.7%
--- + Steroids	0.5%	3.3%	2.3%	0.7%	0.5%	0.6%	0.7%	0.5%	0.4%	0.3%
CyA	0.5%	0.1%	0.3%	0.3%	0.7%	0.6%	0.1%	0.1%	0.1%	0.1%
CyA + MMF/MPA	0.6%	1.2%	0.9%	0.9%	0.5%	1.0%	1.1%	1.5%	1.9%	2.0%
CyA + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.2%	0.0%	0.1%	0.1%	0.0%
CyA + MMF/MPA + Steroids	16.7%	33.7%	37.0%	41.4%	39.2%	31.7%	24.2%	19.9%	15.5%	12.6%
CyA + OtherAntimet	0.5%	0.4%	0.0%	0.3%	0.2%	0.1%	0.0%	0.0%	0.0%	0.0%
CyA + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
CyA + OtherAntimet + Steroids	30.7%	11.4%	5.8%	5.6%	5.4%	3.0%	1.7%	0.9%	0.5%	0.4%
CyA + Siro/Evero	0.2%	0.1%	0.0%	0.0%	0.2%	0.3%	0.2%	0.7%	1.0%	0.9%
CyA + Siro/Evero + Steroids	0.0%	0.9%	1.5%	2.5%	2.1%	4.0%	3.3%	2.7%	2.0%	1.5%
CyA + Steroids	11.9%	11.5%	10.5%	7.8%	6.6%	4.3%	2.5%	1.6%	1.5%	1.2%
Tac	0.5%	0.6%	0.8%	0.5%	0.5%	0.5%	1.1%	1.8%	2.1%	2.2%
Tac + MMF/MPA	0.2%	1.2%	2.3%	1.4%	1.7%	1.8%	2.8%	5.7%	8.2%	14.1%
Tac + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.3%	0.3%	0.4%	0.8%	0.7%
Tac + MMF/MPA + Steroids	4.9%	12.5%	17.6%	24.5%	28.3%	31.5%	37.5%	39.8%	42.6%	42.9%
Tac + OtherAntimet	0.0%	0.1%	0.0%	0.1%	0.1%	0.2%	0.1%	0.1%	0.1%	0.2%
Tac + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Tac + OtherAntimet + Steroids	2.5%	2.5%	2.5%	2.6%	2.2%	1.9%	2.0%	1.5%	1.3%	1.0%
Tac + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.1%	0.7%	1.5%	2.2%	2.4%	2.1%
Tac + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.1%	1.0%	4.7%	7.1%	5.9%	5.1%	3.5%
Tac + Steroids	5.3%	7.2%	6.2%	6.3%	5.9%	6.3%	5.8%	5.7%	4.8%	3.9%
Other Regimen	17.6%	10.1%	9.4%	2.8%	2.8%	4.4%	5.5%	6.4%	6.5%	7.2%

Source: OPTN/SRTR Data as of May 1, 2006.

Notes:

Percentages are calculated based on the total number of patients with maintenance use at one year following transplantation. They sum to 100 percent for each column.

---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.

MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.

"Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.