Table 7.6f
Immunosuppression Use for Maintenance by Regimen
One Year Following Transplantation, 1995 to 2004
Recipients with Pancreas After Kidney (PAK) Transplants

	Transplant Year									
	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
Transplants	67	113	130	156	220	304	305	375	344	420
Functioning Graft at 1 Year PostTx	41	71	89	110	168	219	246	271	260	320
With Maintenance Use Recorded at 1 Year PostTx	2	3	19	70	164	211	243	266	252	317
--- + MMF/MPA	50.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.4%	1.6%	3.2%
--- + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.8%	0.3%
--- + MMF/MPA + Steroids	0.0%	0.0%	0.0%	0.0%	0.6%	0.5%	0.4%	0.4%	0.0%	0.3%
--- + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.3%
--- + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.0%	0.6%	0.5%	0.8%	0.4%	0.4%	0.0%
--- + Steroids	0.0%	0.0%	0.0%	0.0%	0.0%	0.5%	0.0%	0.0%	0.4%	0.0%
CyA	50.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
CyA + MMF/MPA	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.4%	0.8%	0.8%	0.9%
CyA + MMF/MPA + Steroids	0.0%	33.3%	10.5%	22.9%	9.1%	8.1%	8.6%	9.8%	5.2%	3.8%
CyA + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.4%	0.0%	0.0%	0.0%
CyA + OtherAntimet + Steroids	0.0%	0.0%	0.0%	1.4%	0.6%	0.5%	0.0%	0.4%	0.0%	0.0%
CyA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.6%
CyA + Siro/Evero + Steroids	0.0%	0.0%	0.0%	1.4%	1.8%	2.8%	1.2%	1.9%	0.8%	0.6%
CyA + Steroids	0.0%	0.0%	0.0%	1.4%	0.0%	0.9%	0.4%	0.4%	1.2%	0.0%
Tac	0.0%	0.0%	0.0%	0.0%	0.0%	0.9%	0.0%	0.8%	1.6%	5.4%
Tac + MMF/MPA	0.0%	0.0%	0.0%	0.0%	1.2%	2.8%	4.1%	3.4%	7.5%	8.5%
Tac + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.4%	1.2%	1.3%
Tac + MMF/MPA + Steroids	0.0%	66.7%	73.7%	62.9%	73.2%	54.5%	55.1%	53.0%	46.0%	50.5%
Tac + OtherAntimet	0.0%	0.0%	0.0%	1.4%	0.0%	0.9%	0.0%	0.0%	0.8%	0.0%
Tac + OtherAntimet + Steroids	0.0%	0.0%	10.5%	4.3%	4.3%	2.4%	0.8%	0.4%	1.6%	0.9%
Tac + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.9%	4.1%	7.5%	4.4%	7.3%
Tac + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.0%	3.7%	14.7%	15.2%	10.9%	9.9%	7.3%
Tac + Steroids	0.0%	0.0%	5.3%	1.4%	3.7%	4.7%	4.1%	4.1%	3.2%	2.2%
Other Regimen	0.0%	0.0%	0.0%	2.9%	1.2%	4.3%	4.1%	5.3%	12.7%	6.6%

Source: OPTN/SRTR Data as of May 1, 2006.

Notes:

Percentages are calculated based on the total number of patients with maintenance use at one year following transplantation. They sum to 100 percent for each column.

---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.

MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.

"Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.