Table 9.6f
Immunosuppression Use for Maintenance by Regimen
One Year Following Transplantation, 1995 to 2004
Recipients with Liver Transplants

	Transplant Year									
	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
Transplants	3,934	4,082	4,186	4,516	4,751	4,995	5,190	5,331	5,672	6,167
Functioning Graft at 1 Year PostTx	2,977	3,004	3,180	3,484	3,660	3,901	4,094	4,307	4,583	4,982
With Maintenance Use Recorded at 1 Year PostTx	343	424	646	2,258	3,413	3,715	4,026	4,208	4,446	4,861
--- + MMF/MPA	0.0%	0.0%	0.2%	0.1%	0.1%	0.0%	0.1%	0.2%	0.4%	0.3%
--- + MMF/MPA + OtherAntimet	0.0%	0.0%	0.2%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
--- + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.2%	0.4%	1.0%	1.1%	1.3%
--- + MMF/MPA + Steroids	0.0%	0.0%	0.5%	0.3%	0.4%	0.3%	0.3%	0.4%	0.4%	0.4%
--- + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%
--- + OtherAntimet + Steroids	0.0%	0.2%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%
--- + Siro/Evero	0.0%	0.0%	0.0%	0.1%	0.2%	0.8%	1.0%	1.7%	1.5%	2.0%
--- + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.1%	0.1%	0.3%	1.0%	0.9%	0.8%	1.6%
--- + Steroids	0.9%	0.9%	0.5%	1.8%	1.8%	1.6%	0.6%	0.8%	0.5%	0.5%
CyA	0.3%	1.2%	1.1%	3.6%	4.2%	3.1%	2.5%	2.5%	1.9%	1.9%
CyA + MMF/MPA	0.0%	0.7%	0.6%	2.3%	1.6%	1.3%	1.4%	1.6%	1.9%	2.3%
CyA + MMF/MPA + OtherAntimet	0.0%	0.0%	0.5%	0.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
CyA + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%	0.1%
CyA + MMF/MPA + Steroids	3.2%	4.5%	4.8%	4.4%	5.0%	4.0%	3.2%	2.0%	2.1%	1.9%
CyA + OtherAntimet	0.0%	0.2%	0.2%	0.2%	0.2%	0.1%	0.1%	0.0%	0.1%	0.2%
CyA + OtherAntimet + Steroids	12.2%	10.4%	6.5%	2.4%	2.5%	1.7%	0.8%	0.3%	0.2%	0.3%
CyA + Siro/Evero	0.0%	0.0%	0.0%	0.3%	0.1%	0.9%	0.5%	0.3%	0.4%	0.3%
CyA + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.3%	0.3%	0.3%	0.4%	0.3%	0.3%	0.1%
CyA + Steroids	13.1%	15.8%	9.8%	11.1%	10.3%	5.7%	2.9%	2.3%	2.0%	1.7%
Tac	8.7%	8.0%	17.3%	19.4%	20.6%	23.6%	28.9%	31.3%	30.1%	28.3%
Tac + MMF/MPA	0.9%	1.2%	1.9%	4.9%	5.5%	7.1%	7.3%	10.1%	13.7%	15.3%
Tac + MMF/MPA + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Tac + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.2%	0.1%	0.2%	0.3%	0.2%
Tac + MMF/MPA + Steroids	4.4%	7.8%	9.4%	8.8%	10.3%	10.8%	13.2%	12.6%	14.1%	15.4%
Tac + OtherAntimet	0.0%	0.0%	0.5%	0.6%	0.3%	0.5%	0.2%	0.5%	0.3%	0.6%
Tac + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Tac + OtherAntimet + Steroids	4.1%	6.1%	3.4%	2.9%	1.4%	1.1%	1.4%	1.4%	1.0%	0.5%
Tac + Siro/Evero	0.0%	0.0%	0.2%	0.0%	0.4%	2.4%	2.5%	2.3%	2.4%	2.9%
Tac + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.1%	0.6%	3.0%	2.3%	2.1%	1.4%	1.5%
Tac + Steroids	33.8%	35.8%	31.4%	32.8%	30.5%	26.8%	22.7%	22.2%	20.9%	18.1%
Other Regimen	18.4%	7.1%	11.3%	3.2%	3.6%	4.3%	5.8%	2.7%	2.2%	1.8%

Source: OPTN/SRTR Data as of May 1, 2006.

Notes:

Percentages are calculated based on the total number of patients with maintenance use at one year following transplantation. They sum to 100 percent for each column.

---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.

MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.

"Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.