Table 6.6f
Immunosuppression Use for Maintenance by Regimen
One Year Following Transplantation, 1995 to 2004
Recipients with Pancreas Transplant Alone (PTA) Transplants

	Transplant Year									
	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
Transplants	42	52	79	89	139	134	167	180	158	183
Functioning Graft at 1 Year PostTx	23	31	46	64	108	96	122	142	115	139
With Maintenance Use Recorded at 1 Year PostTx	2	3	5	52	107	93	120	141	115	138
--- + MMF/MPA	50.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	1.4%	9.6%	3.6%
--- + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.9%	0.7%
--- + MMF/MPA + Steroids	0.0%	0.0%	0.0%	0.0%	1.9%	1.1%	0.0%	1.4%	0.0%	0.0%
--- + OtherAntimet + Steroids	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.7%
--- + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	1.7%	0.7%	0.9%	2.2%
--- + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.0%	0.0%	2.2%	0.0%	1.4%	0.0%	1.4%
--- + Steroids	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.8%	0.7%	0.0%	0.7%
CyA	0.0%	0.0%	0.0%	0.0%	0.0%	1.1%	0.0%	0.0%	0.0%	0.0%
CyA + MMF/MPA	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.8%	1.4%	0.9%	0.7%
CyA + MMF/MPA + Steroids	0.0%	0.0%	0.0%	5.8%	5.6%	3.2%	4.2%	2.8%	2.6%	1.4%
CyA + OtherAntimet + Steroids	0.0%	0.0%	0.0%	1.9%	0.9%	0.0%	0.0%	0.0%	0.0%	0.0%
CyA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.7%	1.7%	0.0%
CyA + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.0%	0.9%	0.0%	0.0%	0.7%	0.9%	0.7%
CyA + Steroids	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.7%	0.9%	0.0%
Tac	0.0%	0.0%	0.0%	0.0%	1.9%	2.2%	10.8%	5.7%	17.4%	15.2%
Tac + MMF/MPA	0.0%	0.0%	0.0%	0.0%	3.7%	2.2%	2.5%	8.5%	7.0%	12.3%
Tac + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.9%	2.9%
Tac + MMF/MPA + Steroids	0.0%	100.0%	40.0%	75.0%	67.3%	52.7%	55.0%	37.6%	26.1%	22.5%
Tac + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.9%	0.7%
Tac + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.7%
Tac + OtherAntimet + Steroids	0.0%	0.0%	0.0%	7.7%	0.9%	4.3%	0.8%	0.0%	0.0%	2.2%
Tac + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.9%	2.2%	0.8%	9.2%	7.0%	8.0%
Tac + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.0%	8.4%	12.9%	6.7%	11.3%	5.2%	2.9%
Tac + Steroids	0.0%	0.0%	60.0%	9.6%	4.7%	9.7%	15.0%	13.5%	11.3%	12.3%
Other Regimen	50.0%	0.0%	0.0%	0.0%	2.8%	6.5%	0.8%	2.1%	6.1%	8.0%

Source: OPTN/SRTR Data as of May 1, 2006.

Notes:

Percentages are calculated based on the total number of patients with maintenance use at one year following transplantation. They sum to 100 percent for each column.

---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.

MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.

"Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.