Table 10.12
Adjusted Patient Survival, Intestine Transplants
Survival at 3 Months, 1 Year, 3 Years, and 5 Years

	3 Months			1 Year			3 Years			5 Years		
	(Tx 2003 - 2004)			(Tx 2003 - 2004)			(Tx 2001 - 2004)			(Tx 1999 - 2004)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	242	88.9%	2.0%	242	80.2%	2.6%	438	63.0%	2.8%	563	53.5%	3.0%
Age at Tx												
<1 Year	30	93.3%	4.5%	30	66.7%	8.4%	63	61.4%	6.4%	84	52.1%	5.7%
1-5 Years	93	84.9%	3.7%	93	79.6%	4.2%	152	59.6%	5.1%	194	54.6%	4.5%
6-10 Years	14	92.9%	6.6%	14	85.7%	9.0%	27	65.8%	10.0%	36	69.0%	8.2%
11-17 Years	16	93.8%	5.9%	16	93.8%	5.9%	26	76.3%	11.3%	30	61.3%	16.6%
18-34 Years	25	88.0%	6.4%	25	80.0%	7.8%	51	63.8%	7.3%	70	44.2%	8.4%
35-49 Years	40	90.0%	4.7%	40	85.0%	5.6%	75	67.9%	6.8%	99	56.1%	7.5%
50-64 Years	22	95.5%	4.3%	22	86.4%	7.1%	41	55.2%	10.0%	47	48.7%	9.6%
65+ Years	2	50.0%	25.0%	2	0.0%	0.0%	3	33.3%	20.0%	3	33.3%	20.0%
Recipient Ethnicity/Race												
White	176	89.1%	2.3%	176	81.7%	2.9%	310	63.9%	3.4%	392	54.8%	3.7%
African American	25	84.2%	7.1%	25	76.3%	8.3%	56	61.3%	7.1%	75	46.8%	6.7%
Hispanic/Latino	30	100.0%	0.0%	30	86.9%	6.1%	58	63.2%	7.7%	78	57.3%	6.4%
Asian	8	63.4%	15.7%	8	51.1%	16.2%	10	51.8%	14.5%	13	35.4%	15.1%
Other/Multi-race	3	67.5%	21.9%	3	34.5%	20.3%	4	51.0%	20.7%	5	60.2%	19.2%
Recipient Gender												
Female	122	84.4%	3.3%	122	77.9%	3.7%	222	66.0%	3.7%	279	54.5%	4.6%
Male	120	93.4%	2.3%	120	82.6%	3.4%	216	60.0%	4.2%	284	52.2%	3.9%

Source: OPTN/SRTR Data as of May 1, 2006.
Notes:
(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.
Cohorts are transplants performed during the years listed at the top of the table.
Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.
Center volume = Center's yearly transplants performed during the base period, based on intestine transplants.
Multi-organ transplants other than intestine with liver and/or pancreas are excluded.
Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Table 10.12 (Continued)
Adjusted Patient Survival, Intestine Transplants
Survival at 3 Months, 1 Year, 3 Years, and 5 Years

	3 Months			1 Year			3 Years			5 Years		
	(Tx 2003 - 2004)			(Tx 2003 - 2004)			(Tx 2001 - 2004)			(Tx 1999 - 2004)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	242	88.9%	2.0%	242	80.2%	2.6%	438	63.0%	2.8%	563	53.5%	3.0%
Primary Diagnosis												
Short Gut Syndrome	176	88.3%	2.4%	176	78.7%	3.1%	330	63.0%	3.1%	421	54.2%	3.3%
Functional Bowel Problem	45	97.9%	2.0%	45	91.6%	4.0%	75	65.9%	7.1%	101	53.6%	7.4%
Other	20	71.7%	10.7%	20	66.2%	11.3%	32	53.2%	11.5%	40	47.0%	10.7%
Unknown	1	+	+	1	+	+	1	+	+	1	+	+

Source: OPTN/SRTR Data as of May 1, 2006.
Notes:
(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.
Cohorts are transplants performed during the years listed at the top of the table.
Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.
Center volume = Center's yearly transplants performed during the base period, based on intestine transplants.
Multi-organ transplants other than intestine with liver and/or pancreas are excluded.
Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.