Table 5.14b
Unadjusted Patient Survival, Deceased Donor ECD Kidney Transplants
Survival at 3 Months, 1 Year, 3 Years, and 5 Years

	3 Months			1 Year			3 Years			5 Years		
	(Tx 2003 - 2004)			(Tx 2003 - 2004)			(Tx 2001 - 2004)			(Tx 1999 - 2004)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	2,542	96.4%	0.4%	2,542	90.1%	0.6%	4,777	80.7%	0.7%	6,922	69.2%	0.8%
Age at Tx												
6-10 Years	0	-	-	0	-	-	1	100.0%	0.0%	3	66.7%	27.2%
11-17 Years	4	100.0%	0.0%	4	100.0%	0.0%	7	100.0%	0.0%	12	100.0%	0.0%
18-34 Years	88	100.0%	0.0%	88	97.7%	1.6%	220	92.7%	1.9%	383	87.8%	1.9%
35-49 Years	411	98.1%	0.7%	411	95.4%	1.0%	885	89.0%	1.2%	1,397	81.0%	1.4%
50-64 Years	1,246	96.3%	0.5%	1,246	90.3%	0.8%	2,347	79.1%	1.0%	3,394	67.6%	1.1%
65+ Years	793	95.2%	0.8%	793	86.1%	1.2%	1,317	75.4%	1.4%	1,733	54.9%	1.9%
Recipient Ethnicity/Race												
White	1,220	96.6%	0.5%	1,220	89.6%	0.9%	2,347	80.3%	0.9%	3,453	67.0%	1.1%
African American	802	95.9%	0.7%	802	89.7%	1.1%	1,509	79.6%	1.2%	2,150	70.0%	1.3%
Hispanic/Latino	304	97.4%	0.9%	304	91.8%	1.6%	539	83.7%	1.9%	764	73.8%	2.2%
Asian	170	95.3%	1.6%	170	92.4%	2.0%	308	84.1%	2.4%	450	78.1%	2.5%
Other/Multi-race	46	95.7%	3.0%	46	91.3%	4.2%	74	82.5%	5.5%	105	55.9%	7.4%
Recipient Gender												
Female	931	96.5%	0.6%	931	91.3%	0.9%	1,796	82.0%	1.1%	2,642	70.8%	1.2%
Male	1,611	96.3%	0.5%	1,611	89.4%	0.8%	2,981	79.9%	0.9%	4,280	68.3%	1.0%
Recipient Blood Type												
O	1,210	96.8%	0.5%	1,210	91.1%	0.8%	2,214	81.3%	1.0%	3,199	70.5%	1.1%
A	924	96.4%	0.6%	924	89.0%	1.0%	1,762	80.9%	1.1%	2,588	67.5%	1.3%
B	308	95.1%	1.2%	308	88.3%	1.8%	589	76.3%	2.0%	841	68.7%	2.1%
AB	100	95.0%	2.2%	100	94.0%	2.4%	212	84.9%	2.8%	294	72.0%	3.8%

Source: OPTN/SRTR Data as of May 1, 2006.
Notes:
(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.
Cohorts are transplants performed during the years listed at the top of the table.
Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.
Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.
Multi-organ transplants are excluded.
Dialysis needed within first week excludes patients whose graft failed before the end of the week. footnote9

Table 5.14b (Continued)
Unadjusted Patient Survival, Deceased Donor ECD Kidney Transplants
Survival at 3 Months, 1 Year, 3 Years, and 5 Years

	3 Months			1 Year			3 Years			5 Years		
	(Tx 2003 - 2004)			(Tx 2003 - 2004)			(Tx 2001 - 2004)			(Tx 1999 - 2004)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	2,542	96.4%	0.4%	2,542	90.1%	0.6%	4,777	80.7%	0.7%	6,922	69.2%	0.8%
Recipient Residency												
U.S. Resident	2,532	96.4%	0.4%	2,532	90.1%	0.6%	4,752	80.6%	0.7%	6,888	69.1%	0.8%
Nonresident Alien	10	90.0%	9.5%	10	90.0%	9.5%	20	95.0%	4.9%	27	96.3%	3.6%
Unknown	0	-	-	0	-	-	5	80.0%	17.9%	7	85.7%	13.2%
Hospitalized at Tx												
Not Hospitalized	2,502	96.4%	0.4%	2,502	90.2%	0.6%	4,705	80.8%	0.7%	6,804	69.2%	0.8%
Hospitalized	34	97.1%	2.9%	34	82.4%	6.5%	66	72.1%	6.5%	105	71.0%	5.1%
In Intensive Care	6	100.0%	0.0%	6	100.0%	0.0%	6	+	+	12	44.4%	20.3%
Unknown	0	-	-	0	-	-	0	-	-	1	100.0%	0.0%
Life Support at Tx												
Unknown	2,542	96.4%	0.4%	2,542	90.1%	0.6%	4,777	80.7%	0.7%	6,922	69.2%	0.8%
Primary Diagnosis												
Glomerular Diseases	422	96.7%	0.9%	422	92.2%	1.3%	838	86.7%	1.3%	1,293	78.2%	1.5%
Diabetes	800	96.4%	0.7%	800	88.1%	1.1%	1,437	74.6%	1.4%	2,032	58.7%	1.6%
Hypertensive Nephrosclerosis	644	95.7%	0.8%	644	89.4%	1.2%	1,173	80.6%	1.3%	1,592	69.9%	1.6%
Polycystic Kidneys	214	98.1%	0.9%	214	95.3%	1.4%	433	86.3%	2.0%	646	81.8%	2.0%
Tubular and Interstitial Diseases	107	95.3%	2.0%	107	87.9%	3.2%	186	78.0%	3.4%	297	66.0%	3.5%
Renovascular & Other Vascular Diseases	135	95.6%	1.8%	135	87.4%	2.9%	273	78.5%	2.8%	417	67.5%	3.0%
Congenital, Rare Familial, & Metabolic Disorders	18	94.4%	5.4%	18	88.9%	7.4%	45	89.7%	5.0%	62	90.7%	4.0%
Neoplasms	11	100.0%	0.0%	11	90.9%	8.7%	20	77.9%	11.8%	25	79.9%	9.3%
Other	191	97.4%	1.2%	191	93.2%	1.8%	372	85.4%	2.1%	557	67.1%	2.8%
Unknown	0	-	-	0	-	-	0	-	-	1	100.0%	0.0%

Source: OPTN/SRTR Data as of May 1, 2006.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Dialysis needed within first week excludes patients whose graft failed before the end of the week. footnote9

Table 5.14b (Continued)
Unadjusted Patient Survival, Deceased Donor ECD Kidney Transplants
Survival at 3 Months, 1 Year, 3 Years, and 5 Years

	3 Months			1 Year			3 Years			5 Years		
	(Tx 2003 - 2004)			(Tx 2003 - 2004)			(Tx 2001 - 2004)			(Tx 1999 - 2004)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	2,542	96.4%	0.4%	2,542	90.1%	0.6%	4,777	80.7%	0.7%	6,922	69.2%	0.8%
Cold Ischemic Time												
0-11 Hours	429	96.5%	0.9%	429	91.4%	1.4%	707	83.0%	1.7%	900	72.3%	2.3%
12-21 Hours	1,027	97.2%	0.5%	1,027	91.4%	0.9%	1,931	82.2%	1.0%	2,757	71.8%	1.2%
22-31 Hours	557	96.4%	0.8%	557	90.7%	1.2%	1,098	78.8%	1.4%	1,735	67.5%	1.5%
32-41 Hours	139	96.4%	1.6%	139	89.2%	2.6%	293	78.5%	2.8%	474	61.1%	2.9%
42+ Hours	51	96.1%	2.7%	51	86.3%	4.8%	80	83.2%	4.5%	115	74.6%	5.5%
Unknown	339	93.8%	1.3%	339	84.4%	2.0%	668	78.1%	1.8%	941	67.0%	2.1%
Yearly Center Volume												
0-12	7	100.0%	0.0%	7	85.7%	13.2%	34	79.8%	7.5%	60	71.1%	7.0%
13-28	118	95.8%	1.9%	118	91.5%	2.6%	259	76.9%	3.2%	455	68.3%	2.9%
29-47	248	94.4%	1.5%	248	88.3%	2.0%	454	81.0%	2.1%	672	69.2%	2.5%
48-81	515	96.3%	0.8%	515	91.7%	1.2%	1,049	79.6%	1.5%	1,570	69.2%	1.6%
82+	1,654	96.7%	0.4%	1,654	89.8%	0.7%	2,981	81.4%	0.8%	4,165	69.3%	1.0%
Dialysis Needed Within First Week After Tx												
No	1,698	97.3%	0.4%	1,698	91.9%	0.7%	3,137	83.7%	0.8%	4,590	72.8%	0.9%
Yes	844	94.4%	0.8%	844	86.5%	1.2%	1,640	75.1%	1.2%	2,324	61.9%	1.4%
Unknown	0	-	-	0	-	-	0	-	-	8	87.5%	11.7%
Donor Age												
50-64 Years	1,885	96.6%	0.4%	1,885	90.2%	0.7%	3,532	82.0%	0.8%	5,129	71.0%	0.9%
65+ Years	657	95.9%	0.8%	657	89.6%	1.2%	1,245	77.0%	1.4%	1,793	64.0%	1.5%

Source: OPTN/SRTR Data as of May 1, 2006.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Dialysis needed within first week excludes patients whose graft failed before the end of the week. footnote9

Table 5.14b (Continued)
Unadjusted Patient Survival, Deceased Donor ECD Kidney Transplants
Survival at 3 Months, 1 Year, 3 Years, and 5 Years

	3 Months			1 Year			3 Years			5 Years		
	(Tx 2003 - 2004)			(Tx 2003 - 2004)			(Tx 2001 - 2004)			(Tx 1999 - 2004)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	2,542	96.4%	0.4%	2,542	90.1%	0.6%	4,777	80.7%	0.7%	6,922	69.2%	0.8%
A Locus Mismatch												
0	278	97.1%	1.0%	278	93.2%	1.5%	621	84.3%	1.6%	1,021	72.6%	1.7%
1	996	96.6%	0.6%	996	89.9%	1.0%	1,886	81.1%	1.0%	2,687	67.3%	1.3%
2	1,267	96.1%	0.5%	1,267	89.6%	0.9%	2,269	79.3%	1.0%	3,207	69.6%	1.1%
Unknown	1	100.0%	0.0%	1	100.0%	0.0%	1	0.0%	0.0%	7	71.4%	17.1%
B Locus Mismatch												
0	185	97.8%	1.1%	185	92.4%	1.9%	518	83.6%	1.7%	929	71.8%	1.8%
1	660	97.0%	0.7%	660	90.2%	1.2%	1,429	81.7%	1.2%	2,250	71.8%	1.2%
2	1,696	96.0%	0.5%	1,696	89.8%	0.7%	2,829	79.4%	0.9%	3,736	66.3%	1.2%
Unknown	1	100.0%	0.0%	1	100.0%	0.0%	1	0.0%	0.0%	7	71.4%	17.1%
DR Locus Mismatch												
0	315	97.1%	0.9%	315	90.8%	1.6%	846	83.1%	1.4%	1,445	73.7%	1.4%
1	1,040	96.3%	0.6%	1,040	90.2%	0.9%	1,990	80.1%	1.0%	2,952	69.8%	1.1%
2	1,187	96.3%	0.5%	1,187	89.8%	0.9%	1,941	80.3%	1.1%	2,518	64.2%	1.5%
Unknown	0	-	-	0	-	-	0	-	-	7	85.7%	13.2%
Total Mismatch												
0	131	97.7%	1.3%	131	93.9%	2.1%	307	86.0%	2.2%	525	73.8%	2.4%
1	19	94.7%	5.1%	19	89.5%	7.0%	75	85.3%	4.1%	170	70.8%	3.9%
2	85	97.6%	1.6%	85	92.9%	2.8%	282	82.7%	2.4%	490	75.5%	2.2%
3	290	97.6%	0.9%	290	90.0%	1.8%	720	81.3%	1.6%	1,136	70.3%	1.7%
4	636	96.9%	0.7%	636	89.6%	1.2%	1,198	80.2%	1.4%	1,745	69.0%	1.5%
5	888	95.2%	0.7%	888	89.8%	1.0%	1,373	78.9%	1.4%	1,816	66.3%	1.7%
6	492	96.7%	0.8%	492	89.8%	1.4%	821	80.1%	1.7%	1,032	64.3%	2.4%
Unknown	1	100.0%	0.0%	1	100.0%	0.0%	1	0.0%	0.0%	8	75.0%	15.3%

Source: OPTN/SRTR Data as of May 1, 2006.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Dialysis needed within first week excludes patients whose graft failed before the end of the week. footnote9