Table 7.6d
Immunosuppression Use for Maintenance by Regimen
Prior to Discharge, 1996 to 2005
Recipients with Pancreas After Kidney (PAK) Transplants

	Transplant Year									
	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Functioning Graft at Discharge	98	114	139	206	277	287	336	316	381	307
With Immunosuppression Info	94	113	137	203	276	286	335	313	375	299
--- + MMF/MPA	3.2%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	5.8%	4.8%	2.0%
--- + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.3%	1.6%	0.3%
--- + MMF/MPA + Steroids	8.5%	2.7%	0.7%	1.0%	0.0%	0.3%	1.5%	1.6%	1.6%	0.3%
--- + OtherAntimet	1.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
--- + OtherAntimet + Steroids	3.2%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
--- + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.5%	0.3%
--- + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.3%	0.3%	0.3%	0.3%
--- + Steroids	3.2%	0.0%	0.7%	2.0%	1.8%	0.3%	0.3%	1.0%	2.7%	3.3%
CyA	0.0%	0.0%	0.0%	0.0%	0.0%	0.3%	0.3%	0.0%	0.0%	0.0%
CyA + MMF/MPA	4.3%	0.0%	0.0%	0.5%	0.0%	0.7%	0.3%	1.0%	1.3%	2.0%
CyA + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.3%
CyA + MMF/MPA + Steroids	20.2%	16.8%	18.2%	10.3%	10.9%	5.6%	7.2%	5.1%	3.5%	2.3%
CyA + OtherAntimet + Steroids	5.3%	0.0%	0.7%	0.0%	0.0%	0.0%	0.0%	0.3%	0.0%	0.0%
CyA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	1.1%	1.0%
CyA + Siro/Evero + Steroids	0.0%	0.0%	0.7%	0.5%	1.8%	1.4%	2.1%	1.3%	0.8%	0.7%
CyA + Steroids	2.1%	0.0%	1.5%	0.5%	0.0%	0.0%	0.3%	0.3%	0.5%	0.0%
Tac	3.2%	0.0%	0.0%	6.9%	4.0%	3.8%	4.8%	1.3%	2.7%	3.0%
Tac + MMF/MPA	2.1%	2.7%	0.0%	0.5%	0.0%	0.3%	0.9%	6.7%	8.3%	17.1%
Tac + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.3%	1.6%	0.8%	1.7%
Tac + MMF/MPA + Steroids	40.4%	67.3%	65.7%	68.5%	67.4%	62.6%	60.6%	59.7%	55.2%	54.5%
Tac + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.3%
Tac + OtherAntimet + Steroids	1.1%	0.9%	1.5%	2.0%	1.8%	0.7%	0.3%	0.3%	0.3%	0.7%
Tac + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	1.0%	2.4%	2.2%	4.0%	4.3%
Tac + Siro/Evero + Steroids	0.0%	0.0%	0.0%	1.0%	5.4%	15.7%	11.3%	6.1%	4.5%	2.3%
Tac + Steroids	1.1%	3.5%	3.6%	2.5%	4.3%	1.0%	2.4%	1.3%	2.1%	0.3%
Other Regimen	1.1%	6.2%	6.6%	3.9%	2.5%	5.9%	4.8%	3.8%	3.5%	2.7%

Source: OPTN/SRTR Data as of May 1, 2006.
Notes:
Percentages in each column sum to 100 percent.
---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.
MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.
"Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.