Table 8.6b
Immunosuppression Use for Induction by Discharge Regimen, 2001 to 2005
Recipients with Kidney-Pancreas Transplants

	Discharge Regimen (w/ or w/o Steroids Use)										All
	CyA	Tac	CyA+ MMF	Tac+ MMF	CyA+ Aza	Tac+ Aza	Siro+ MMF	CyA+ Siro	Tac+ Siro	Other	
Functioning Graft with Immunosuppression Info at Discharge	5	169	215	2,945	2	14	89	110	412	347	4,308
% of All Regimens	0.1%	3.9%	5.0%	68.4%	0.0%	0.3%	2.1%	2.6%	9.6%	8.1%	100.0%
Induction Drug											
Atgam/NRATG/NRATS	0.0%	0.6%	2.8%	2.9%	50.0%	0.0%	1.1%	0.0%	1.0%	2.3%	2.5%
OKT3	0.0%	0.0%	0.9%	1.3%	0.0%	0.0%	2.2%	0.9%	1.5%	2.3%	1.3%
Thymoglobulin	60.0%	31.4%	33.0%	40.0%	0.0%	14.3%	13.5%	79.1%	53.9%	67.4%	43.2%
Zenapax	20.0%	4.1%	8.4%	11.9%	0.0%	14.3%	2.2%	1.8%	13.3%	12.4%	11.1%
Simulect	0.0%	5.9%	46.5%	20.5%	0.0%	42.9%	11.2%	0.9%	6.6%	21.3%	19.3%
Campath	0.0%	21.3%	0.5%	5.0%	0.0%	0.0%	64.0%	0.9%	10.7%	20.2%	8.2%
No Induction Drugs Recorded	40.0%	39.1%	13.0%	22.9%	50.0%	28.6%	11.2%	19.1%	26.5%	13.3%	22.3%

Source: OPTN/SRTR Data as of May 1, 2006.
Notes:
Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.
CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.