Table 10.6d
Immunosuppression Use for Maintenance by Regimen
Prior to Discharge, 1996 to 2005
Recipients with Intestine Transplants

	Transplant Year									
	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Functioning Graft at Discharge	35	45	47	49	61	88	89	103	130	153
With Immunosuppression Info	24	44	47	49	61	88	89	103	127	149
--- + MMF/MPA	0.0%	0.0%	0.0%	0.0%	1.6%	0.0%	0.0%	0.0%	0.0%	0.0%
--- + MMF/MPA + Steroids	8.3%	0.0%	2.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.7%
--- + OtherAntimet + Steroids	0.0%	0.0%	0.0%	0.0%	1.6%	0.0%	0.0%	0.0%	0.0%	0.0%
--- + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.0%	1.6%	1.1%	0.0%	0.0%	1.6%	0.7%
--- + Steroids	0.0%	2.3%	0.0%	0.0%	0.0%	1.1%	2.2%	3.9%	0.8%	2.0%
CyA + Steroids	0.0%	2.3%	6.4%	4.1%	0.0%	0.0%	0.0%	1.0%	0.0%	0.0%
Tac	0.0%	0.0%	0.0%	4.1%	1.6%	18.2%	32.6%	30.1%	22.8%	25.5%
Tac + MMF/MPA	0.0%	2.3%	0.0%	2.0%	0.0%	0.0%	1.1%	0.0%	0.0%	0.7%
Tac + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	1.0%	0.8%	0.0%
Tac + MMF/MPA + Steroids	25.0%	34.1%	27.7%	14.3%	13.1%	3.4%	7.9%	6.8%	8.7%	10.1%
Tac + OtherAntimet + Steroids	20.8%	9.1%	2.1%	6.1%	0.0%	6.8%	0.0%	0.0%	0.0%	0.0%
Tac + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	2.3%	0.0%	0.0%	1.6%	1.3%
Tac + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.0%	34.4%	18.2%	15.7%	3.9%	7.9%	10.1%
Tac + Steroids	41.7%	43.2%	55.3%	67.3%	41.0%	48.9%	34.8%	50.5%	55.9%	49.0%
Other Regimen	4.2%	6.8%	6.4%	2.0%	4.9%	0.0%	5.6%	2.9%	0.0%	0.0%

Source: OPTN/SRTR Data as of May 1, 2006.
Notes:
Percentages in each column sum to 100 percent.
---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.
MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.
"Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.