Table 13.6b
Immunosuppression Use for Induction by Discharge Regimen, 2001 to 2005
Recipients with Heart-Lung Transplants

	Discharge Regimen (w/ or w/o Steroids Use)								
	CyA	Tac	CyA+ MMF	Tac+ MMF	CyA+ Aza	Tac+ Aza	Tac+ Siro	Other	All
Functioning Graft with Immunosuppression Info at Discharge	1	9	23	32	33	16	1	6	121
% of All Regimens	0.8%	7.4%	19.0%	26.4%	27.3%	13.2%	0.8%	5.0%	100.0%
Induction Drug									
Atgam/NRATG/NRATS	0.0%	0.0%	13.0%	6.3%	39.4%	0.0%	0.0%	66.7%	18.2%
OKT3	0.0%	0.0%	4.3%	3.1%	9.1%	0.0%	0.0%	0.0%	4.1%
Thymoglobulin	0.0%	0.0%	4.3%	18.8%	0.0%	18.8%	100.0%	33.3%	10.7%
Zenapax	0.0%	0.0%	17.4%	9.4%	12.1%	31.3%	0.0%	66.7%	16.5%
Simulect	0.0%	0.0%	39.1%	9.4%	15.2%	25.0%	0.0%	166.7%	25.6%
Campath	0.0%	11.1%	0.0%	9.4%	0.0%	0.0%	0.0%	16.7%	4.1%
No Induction Drugs Recorded	100.0%	88.9%	26.1%	43.8%	30.3%	25.0%	0.0%	0.0%	35.5%

Source: OPTN/SRTR Data as of May 1, 2006.
Notes:
Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.
CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.