Table 2.16
Deceased Donor Organ Utilization, 1996 to 2005, Intestine

	Year									
	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total # Donors of Any Organ	5,418	5,479	5,793	5,824	5,985	6,080	6,190	6,457	7,150	7,593
# Intestines Recovered	48	72	78	95	87	115	112	122	166	184
ORPD (1)	0.01	0.01	0.01	0.02	0.01	0.02	0.02	0.02	0.02	0.02
# Intestines Transplanted	42	65	68	71	79	111	106	111	146	171
OTPD (2)	0.01	0.01	0.01	0.01	0.01	0.02	0.02	0.02	0.02	0.02
% Transplated	87.5%	90.3%	87.2%	74.7%	90.8%	96.5%	94.6%	91.0%	88.0%	92.9%
Donor with Brain Death(DBD)										
Total # Donors of Any Organ	5,362	5,410	5,726	5,748	5,880	5,935	6,033	6,237	6,829	7,134
# Intestines Recovered	48	72	78	95	87	115	112	121	166	184
ORPD (1)	0.01	0.01	0.01	0.02	0.01	0.02	0.02	0.02	0.02	0.03
# Intestines Transplanted	42	65	68	71	79	111	106	111	146	171
OTPD (2)	0.01	0.01	0.01	0.01	0.01	0.02	0.02	0.02	0.02	0.02
% Transplated	87.5%	90.3%	87.2%	74.7%	90.8%	96.5%	94.6%	91.7%	88.0%	92.9%
Donors After Cardiac Death(DCD)										
Total # Donors of Any Organ	56	69	67	76	105	145	157	220	321	459
# Intestines Recovered	-	-	-	-	-	-	-	1	-	-
ORPD (1)	-	-	-	-	-	-	-	0.00	-	-
# Intestines Transplanted	-	-	-	-	-	-	-	0	-	-
OTPD (2)	-	-	-	-	-	-	-	0.00	-	-
% Transplated	-	-	-	-	-	-	-	0.0%	-	-
Intestine Donors	48	72	78	95	87	115	112	122	166	184
Intestine Donors (%)	0.9%	1.3%	1.3%	1.6%	1.5%	1.9%	1.8%	1.9%	2.3%	2.4%
Intestine Donors, Other Organs Recovered										
Intestine Only	-	1	-	-	1	-	2	-	-	1
Intestine & Kidney or Pancreas Only	21	23	26	32	28	37	39	29	45	46
Intestine & Liver Only	1	5	6	4	4	3	3	5	7	5

(Continued)

Source: OPTN/SRTR Data as of May 1, 2006.
Notes:
(-) = None in category.
(1) ORPD: Organs Recovered Per Donor of any organ; (2) OTPD: Organs Transplanted Per Donor of any organ. Organs divided into segments (liver, lung, pancreas, intestine) are counted only once for that organ.
Includes organs recovered for transplant but not transplanted, as well as organs transplanted.
Donors are divided into two mutually exclusive and complete categories. DCDs include patients whose hearts have irreversibly stopped beating, and may also have been known as Non-Heart Beating Donors. DBDs have suffered brain death.

Table 2.16 (Continued)
Deceased Donor Organ Utilization, 1996 to 2005, Intestine

	Year									
	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Intestine Donors, Other Organs Recovered										
Intestine & Thoracic Only	3	5	10	9	9	9	7	10	14	11
Intestine & Other Combinations	23	38	36	50	45	66	61	78	100	121
Intestine Donors, Other Organs Recovered (%)										
Intestine Only	-	1.4%	-	-	1.1%	-	1.8%	-	-	0.5%
Intestine & Kidney or Pancreas Only	43.8%	31.9%	33.3%	33.7%	32.2%	32.2%	34.8%	23.8%	27.1%	25.0%
Intestine & Liver Only	2.1%	6.9%	7.7%	4.2%	4.6%	2.6%	2.7%	4.1%	4.2%	2.7%
Intestine & Thoracic Only	6.3%	6.9%	12.8%	9.5%	10.3%	7.8%	6.3%	8.2%	8.4%	6.0%
Intestine & Other Combinations	47.9%	52.8%	46.2%	52.6%	51.7%	57.4%	54.5%	63.9%	60.2%	65.8%

Source: OPTN/SRTR Data as of May 1, 2006.
Notes:
(-) = None in category.
(1) ORPD: Organs Recovered Per Donor of any organ; (2) OTPD: Organs Transplanted Per Donor of any organ. Organs divided into segments (liver, lung, pancreas, intestine) are counted only once for that organ.
Includes organs recovered for transplant but not transplanted, as well as organs transplanted.
Donors are divided into two mutually exclusive and complete categories. DCDs include patients whose hearts have irreversibly stopped beating, and may also have been known as Non-Heart Beating Donors. DBDs have suffered brain death.