Table 9.6b
Immunosuppression Use for Induction by Discharge Regimen, 2001 to 2005
Recipients with Liver Transplants

	Discharge Regimen (w/ or w/o Steroids Use)										All
	CyA	Tac	CyA+ MMF	Tac+ MMF	CyA+ Aza	Tac+ Aza	Siro+ MMF	CyA+ Siro	Tac+ Siro	Other	
Functioning Graft with Immunosuppression Info at Discharge	501	9,407	1,132	11,678	217	344	249	150	616	1,461	25,755
% of All Regimens	1.9%	36.5%	4.4%	45.3%	0.8%	1.3%	1.0%	0.6%	2.4%	5.7%	100.0%
Induction Drug											
Atgam/NRATG/NRATS	0.2%	0.4%	0.7%	0.2%	0.0%	0.0%	0.0%	0.0%	0.5%	1.8%	0.4%
OKT3	0.4%	0.6%	0.4%	0.3%	0.0%	0.6%	0.8%	0.0%	0.8%	1.5%	0.5%
Thymoglobulin	3.8%	7.6%	4.1%	4.0%	10.6%	7.3%	2.8%	1.3%	8.1%	15.8%	6.1%
Zenapax	5.2%	2.8%	6.5%	5.7%	7.8%	16.0%	8.0%	8.0%	3.9%	13.2%	5.3%
Simulect	8.0%	4.9%	20.7%	6.0%	2.3%	43.6%	12.9%	5.3%	3.9%	18.8%	7.5%
Campath	0.2%	1.6%	0.1%	0.5%	0.0%	0.0%	0.0%	0.0%	0.5%	1.0%	0.9%
No Induction Drugs Recorded	82.6%	82.4%	67.8%	83.5%	79.3%	32.6%	75.5%	85.3%	82.8%	80.2%	81.4%

Source: OPTN/SRTR Data as of May 1, 2006.
Notes:
Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.
CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.