

Table 2.15
Deceased Donor Organ Utilization, 2000 to 2009, Liver

	Year									
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Total # Donors of Any Organ	5,985	6,080	6,190	6,457	7,150	7,593	8,019	8,086	7,990	8,022
# Livers Recovered	4,997	5,106	5,294	5,682	6,319	6,693	7,017	6,937	6,752	6,740
ORPD (1)	0.83	0.84	0.86	0.88	0.88	0.88	0.88	0.86	0.85	0.84
# Livers Transplanted	4,505	4,598	4,893	5,263	5,777	6,041	6,309	6,154	5,999	6,016
OTPD (2)	0.75	0.76	0.79	0.82	0.81	0.80	0.79	0.76	0.75	0.75
% Transplanted	90.2%	90.1%	92.4%	92.6%	91.4%	90.3%	89.9%	88.7%	88.8%	89.3%
Donor with Brain Death(DBD)										
Total # Donors of Any Organ	5,868	5,913	6,001	6,189	6,757	7,029	7,375	7,295	7,142	7,102
# Livers Recovered	4,944	5,016	5,193	5,527	6,076	6,303	6,609	6,494	6,344	6,305
ORPD (1)	0.84	0.85	0.87	0.89	0.90	0.90	0.90	0.89	0.89	0.89
# Livers Transplanted	4,466	4,531	4,814	5,151	5,593	5,769	6,020	5,848	5,723	5,727
OTPD (2)	0.76	0.77	0.80	0.83	0.83	0.82	0.82	0.80	0.80	0.81
% Transplanted	90.3%	90.3%	92.7%	93.2%	92.1%	91.5%	91.1%	90.1%	90.2%	90.8%
Donors After Cardiac Death(DCD)										
Total # Donors of Any Organ	117	167	189	268	393	564	644	791	848	920
# Livers Recovered	53	90	101	155	243	390	408	443	408	435
ORPD (1)	0.45	0.54	0.53	0.58	0.62	0.69	0.63	0.56	0.48	0.47
# Livers Transplanted	39	67	79	112	184	272	289	306	276	289
OTPD (2)	0.33	0.40	0.42	0.42	0.47	0.48	0.45	0.39	0.33	0.31
% Transplanted	73.6%	74.4%	78.2%	72.3%	75.7%	69.7%	70.8%	69.1%	67.6%	66.4%
Liver Donors	4,997	5,106	5,294	5,682	6,319	6,693	7,017	6,937	6,752	6,740
Liver Donors (%)	83.5%	84.0%	85.5%	88.0%	88.4%	88.1%	87.5%	85.8%	84.5%	84.0%
Liver Donors, Other Organs Recovered										
Liver Only	321	377	382	532	655	699	679	665	623	614
Liver & Kidney or Pancreas Only	2,423	2,412	2,606	2,881	3,303	3,421	3,689	3,588	3,493	3,366
Liver & Intestine Only	5	4	3	5	6	5	2	5	-	1
Liver & Thoracic Only	73	70	71	84	80	102	91	85	88	91
Liver & Other Combinations	2,175	2,243	2,232	2,180	2,275	2,466	2,556	2,594	2,548	2,668
Liver Donors, Other Organs Recovered (%)										
Liver Only	6.4%	7.4%	7.2%	9.4%	10.4%	10.4%	9.7%	9.6%	9.2%	9.1%

(Continued)

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(1) ORPD: Organs Recovered Per Donor of any organ; (2) OTPD: Organs Transplanted Per Donor of any organ. Organs divided into segments (liver, lung, pancreas, intestine) are counted only once for that organ.

Includes organs recovered for transplant but not transplanted, as well as organs transplanted.

Donors are divided into two mutually exclusive and complete categories. DCDs include patients whose hearts have irreversibly stopped beating, and may also have been known as Non-Heart Beating Donors. DBDs have suffered brain death.

Table 2.15 (Continued)
Deceased Donor Organ Utilization, 2000 to 2009, Liver

	Year									
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Liver Donors, Other Organs Recovered (%)										
Liver & Kidney or Pancreas Only	48.5%	47.2%	49.2%	50.7%	52.3%	51.1%	52.6%	51.7%	51.7%	49.9%
Liver & Intestine Only	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	0.0%	0.1%	-	0.0%
Liver & Thoracic Only	1.5%	1.4%	1.3%	1.5%	1.3%	1.5%	1.3%	1.2%	1.3%	1.4%
Liver & Other Combinations	43.5%	43.9%	42.2%	38.4%	36.0%	36.8%	36.4%	37.4%	37.7%	39.6%

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(1) ORPD: Organs Recovered Per Donor of any organ; (2) OTPD: Organs Transplanted Per Donor of any organ. Organs divided into segments (liver, lung, pancreas, intestine) are counted only once for that organ.

Includes organs recovered for transplant but not transplanted, as well as organs transplanted.

Donors are divided into two mutually exclusive and complete categories. DCDs include patients whose hearts have irreversibly stopped beating, and may also have been known as Non-Heart Beating Donors. DBDs have suffered brain death.