

**Table 2.16**  
**Deceased Donor Organ Utilization, 2000 to 2009, Intestine**

	Year									
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
<b>Total # Donors of Any Organ</b>	5,985	6,080	6,190	6,457	7,150	7,593	8,019	8,086	7,990	8,022
# Intestines Recovered	87	116	112	123	166	184	185	205	197	187
ORPD (1)	0.01	0.02	0.02	0.02	0.02	0.02	0.02	0.03	0.02	0.02
# Intestines Transplanted	79	112	106	112	146	171	171	197	185	178
OTPD (2)	0.01	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
% Transplanted	90.8%	96.6%	94.6%	91.1%	88.0%	92.9%	92.4%	96.1%	93.9%	95.2%
<b>Donor with Brain Death (DBD)</b>										
Total # Donors of Any Organ	5,868	5,913	6,001	6,189	6,757	7,029	7,375	7,295	7,142	7,102
# Intestines Recovered	87	116	112	122	166	184	185	205	197	187
ORPD (1)	0.01	0.02	0.02	0.02	0.02	0.03	0.03	0.03	0.03	0.03
# Intestines Transplanted	79	112	106	112	146	171	171	197	185	178
OTPD (2)	0.01	0.02	0.02	0.02	0.02	0.02	0.02	0.03	0.03	0.03
% Transplanted	90.8%	96.6%	94.6%	91.8%	88.0%	92.9%	92.4%	96.1%	93.9%	95.2%
<b>Donors After Cardiac Death (DCD)</b>										
Total # Donors of Any Organ	117	167	189	268	393	564	644	791	848	920
# Intestines Recovered	-	-	-	1	-	-	-	-	-	-
ORPD (1)	-	-	-	0.00	-	-	-	-	-	-
# Intestines Transplanted	-	-	-	0	-	-	-	-	-	-
OTPD (2)	-	-	-	0.00	-	-	-	-	-	-
% Transplanted	-	-	-	0.0%	-	-	-	-	-	-
<b>Intestine Donors</b>	87	116	112	123	166	184	185	205	197	187
<b>Intestine Donors (%)</b>	1.5%	1.9%	1.8%	1.9%	2.3%	2.4%	2.3%	2.5%	2.5%	2.3%
<b>Intestine Donors, Other Organs Recovered</b>										
Intestine Only	1	-	2	-	-	1	-	-	-	-
Intestine & Kidney or Pancreas Only	27	37	39	29	46	46	49	54	40	48
Intestine & Liver Only	5	4	3	5	6	5	2	5	-	1

(Continued)

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(1) ORPD: Organs Recovered Per Donor of any organ; (2) OTPD: Organs Transplanted Per Donor of any organ. Organs divided into segments (liver, lung, pancreas, intestine) are counted only once for that organ.

Includes organs recovered for transplant but not transplanted, as well as organs transplanted.

Donors are divided into two mutually exclusive and complete categories. DCDs include patients whose hearts have irreversibly stopped beating, and may also have been known as Non-Heart Beating Donors. DBDs have suffered brain death.

**Table 2.16 (Continued)**  
**Deceased Donor Organ Utilization, 2000 to 2009, Intestine**

	Year									
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
<b>Intestine Donors, Other Organs Recovered</b>										
Intestine & Thoracic Only	9	10	7	11	13	13	13	7	5	3
Intestine & Other Combinations	45	65	61	78	101	119	121	139	152	135
<b>Intestine Donors, Other Organs Recovered (%)</b>										
Intestine Only	1.1%	-	1.8%	-	-	0.5%	-	-	-	-
Intestine & Kidney or Pancreas Only	31.0%	31.9%	34.8%	23.6%	27.7%	25.0%	26.5%	26.3%	20.3%	25.7%
Intestine & Liver Only	5.7%	3.4%	2.7%	4.1%	3.6%	2.7%	1.1%	2.4%	-	0.5%
Intestine & Thoracic Only	10.3%	8.6%	6.3%	8.9%	7.8%	7.1%	7.0%	3.4%	2.5%	1.6%
Intestine & Other Combinations	51.7%	56.0%	54.5%	63.4%	60.8%	64.7%	65.4%	67.8%	77.2%	72.2%

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(1) ORPD: Organs Recovered Per Donor of any organ; (2) OTPD: Organs Transplanted Per Donor of any organ. Organs divided into segments (liver, lung, pancreas, intestine) are counted only once for that organ.

Includes organs recovered for transplant but not transplanted, as well as organs transplanted.

Donors are divided into two mutually exclusive and complete categories. DCDs include patients whose hearts have irreversibly stopped beating, and may also have been known as Non-Heart Beating Donors. DBDs have suffered brain death.