

Table 5.7b
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

Recipient Age at Tx	Year of Transplant								
	2000	2001	2002	2003	2004	2005	2006	2007	2008
Total									
Patients	1,170	1,188	1,270	1,442	1,524	1,727	1,756	1,792	1,877
Deaths	110	124	118	140	143	167	152	122	124
Rate	99.3	111.7	99.0	103.3	99.3	102.4	91.6	70.9	68.7
6-11 Years									
Patients	1	1	-	-	2	-	-	-	-
Deaths	1	0	-	-	0	-	-	-	-
Rate	*	*	-	-	*	-	-	-	-
12-17 Years									
Patients	3	2	1	1	1	-	1	-	-
Deaths	0	0	0	0	0	-	0	-	-
Rate	*	*	*	*	*	-	*	-	-
18-34 Years									
Patients	107	79	89	63	70	55	50	39	30
Deaths	2	2	3	1	5	3	1	0	0
Rate	18.8	25.6	34.4	15.9	73.7	56.5	20.4	0.0	0.0
35-49 Years									
Patients	298	272	279	259	277	281	218	188	220
Deaths	12	14	15	15	11	12	8	8	7
Rate	41.0	53.3	55.5	60.0	40.9	43.8	37.7	43.7	32.5
50-64 Years									
Patients	532	585	609	709	716	790	870	938	952
Deaths	61	59	60	68	68	74	66	62	53
Rate	122.8	107.6	105.5	101.8	100.5	99.0	79.5	68.4	57.5
65+ Years									
Patients	229	249	292	410	458	601	617	627	675
Deaths	34	49	40	56	59	78	77	52	64
Rate	162.7	224.2	151.0	149.8	139.3	140.4	135.7	87.9	100.5

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.7b (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

Ethnicity/Race	Year of Transplant								
	2000	2001	2002	2003	2004	2005	2006	2007	2008
Total									
Patients	1,170	1,188	1,270	1,442	1,524	1,727	1,756	1,792	1,877
Deaths	110	124	118	140	143	167	152	122	124
Rate	99.3	111.7	99.0	103.3	99.3	102.4	91.6	70.9	68.7
White									
Patients	616	631	637	712	713	847	876	854	844
Deaths	63	68	62	69	74	70	83	69	69
Rate	108.4	115.4	104.4	103.6	109.8	86.8	100.6	84.6	85.6
African American									
Patients	352	352	410	480	447	525	569	567	599
Deaths	31	34	38	54	37	51	49	34	31
Rate	93.1	102.9	98.7	120.2	87.7	102.9	91.7	62.3	53.5
Other/Multi-race									
Patients	14	16	13	15	37	31	20	24	30
Deaths	2	2	1	1	3	5	1	2	0
Rate	151.7	131.1	77.9	69.2	86.8	182.6	52.5	87.3	0.0
Asian									
Patients	77	68	76	91	112	121	123	128	132
Deaths	2	10	10	8	8	13	5	8	6
Rate	26.2	164.5	143.8	93.6	75.3	114.5	41.5	65.5	46.8
Hispanic/Latino									
Patients	111	121	134	144	215	203	168	219	272
Deaths	12	10	7	8	21	28	14	9	18
Rate	115.9	87.5	53.5	57.2	103.2	149.5	87.4	42.1	69.3

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.7b (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

Gender	Year of Transplant								
	2000	2001	2002	2003	2004	2005	2006	2007	2008
Total									
Patients	1,170	1,188	1,270	1,442	1,524	1,727	1,756	1,792	1,877
Deaths	110	124	118	140	143	167	152	122	124
Rate	99.3	111.7	99.0	103.3	99.3	102.4	91.6	70.9	68.7
Female									
Patients	484	465	499	518	565	625	646	660	678
Deaths	36	43	41	53	44	41	51	36	37
Rate	77.7	98.1	87.0	109.5	81.7	68.2	83.2	56.6	56.5
Male									
Patients	686	723	771	924	959	1,102	1,110	1,132	1,199
Deaths	74	81	77	87	99	126	101	86	87
Rate	114.9	120.6	106.8	99.9	109.8	122.5	96.5	79.3	75.7

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.7b (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

Blood Type	Year of Transplant								
	2000	2001	2002	2003	2004	2005	2006	2007	2008
Total									
Patients	1,170	1,188	1,270	1,442	1,524	1,727	1,756	1,792	1,877
Deaths	110	124	118	140	143	167	152	122	124
Rate	99.3	111.7	99.0	103.3	99.3	102.4	91.6	70.9	68.7
A									
Patients	459	460	460	530	555	613	638	616	645
Deaths	52	47	39	51	62	61	55	49	46
Rate	120.9	108.9	89.9	102.4	119.4	106.0	90.8	83.8	74.5
AB									
Patients	43	70	56	66	65	86	81	82	70
Deaths	4	8	5	5	4	5	12	7	3
Rate	95.6	121.6	95.5	80.4	64.2	60.1	163.1	90.7	43.6
B									
Patients	118	151	161	168	189	216	206	202	228
Deaths	4	20	24	22	16	23	16	16	16
Rate	34.6	144.6	164.8	142.2	88.9	112.8	81.9	83.2	73.6
O									
Patients	550	507	593	678	715	812	831	892	934
Deaths	50	49	50	62	61	78	69	50	59
Rate	96.2	103.3	89.2	96.8	89.9	101.6	87.9	57.8	65.5

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.7b (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

Residency	Year of Transplant								
	2000	2001	2002	2003	2004	2005	2006	2007	2008
Total									
Patients	1,170	1,188	1,270	1,442	1,524	1,727	1,756	1,792	1,877
Deaths	110	124	118	140	143	167	152	122	124
Rate	99.3	111.7	99.0	103.3	99.3	102.4	91.6	70.9	68.7
U.S. Resident									
Patients	1,163	1,178	1,262	1,437	1,519	1,721	1,746	1,788	1,869
Deaths	110	124	117	140	143	165	152	122	124
Rate	100.0	112.7	98.7	103.7	99.6	101.5	92.1	71.1	69.0
Nonresident Alien									
Patients	3	7	5	5	5	5	9	4	8
Deaths	0	0	0	0	0	1	0	0	0
Rate	*	*	*	*	*	*	*	*	*
Unknown									
Patients	4	3	3	-	-	1	1	-	-
Deaths	0	0	1	-	-	1	0	-	-
Rate	*	*	*	-	-	*	*	-	-

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.7b (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

Previous Tx Same Organ	Year of Transplant								
	2000	2001	2002	2003	2004	2005	2006	2007	2008
Total									
Patients	1,170	1,188	1,270	1,442	1,524	1,727	1,756	1,792	1,877
Deaths	110	124	118	140	143	167	152	122	124
Rate	99.3	111.7	99.0	103.3	99.3	102.4	91.6	70.9	68.7
No									
Patients	1,044	1,084	1,184	1,324	1,399	1,592	1,624	1,666	1,768
Deaths	103	119	109	131	132	154	141	111	120
Rate	104.6	117.9	98.0	105.4	99.8	102.5	91.9	69.3	70.7
Yes									
Patients	126	104	86	118	125	135	132	126	109
Deaths	7	5	9	9	11	13	11	11	4
Rate	57.1	49.4	112.7	80.4	93.2	102.0	87.4	92.9	37.5

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.7b (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

Hospitalized at Tx	Year of Transplant								
	2000	2001	2002	2003	2004	2005	2006	2007	2008
Total									
Patients	1,170	1,188	1,270	1,442	1,524	1,727	1,756	1,792	1,877
Deaths	110	124	118	140	143	167	152	122	124
Rate	99.3	111.7	99.0	103.3	99.3	102.4	91.6	70.9	68.7
Not Hospitalized									
Patients	1,151	1,164	1,254	1,421	1,497	1,703	1,731	1,765	1,853
Deaths	106	121	115	135	139	162	149	119	123
Rate	97.1	111.2	97.6	100.9	98.2	100.6	91.1	70.2	69.1
Hospitalized									
Patients	15	23	16	20	23	15	20	25	22
Deaths	2	3	3	5	4	4	1	2	1
Rate	146.9	144.5	221.4	297.5	193.7	334.2	52.4	84.0	46.3
In Intensive Care									
Patients	3	1	-	1	4	9	5	2	2
Deaths	2	0	-	0	0	1	2	1	0
Rate	*	*	-	*	*	*	*	*	*
Unknown									
Patients	1	-	-	-	-	-	-	-	-
Deaths	0	-	-	-	-	-	-	-	-
Rate	*	-	-	-	-	-	-	-	-

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.7b (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

Primary Diagnosis	Year of Transplant								
	2000	2001	2002	2003	2004	2005	2006	2007	2008
Total									
Patients	1,170	1,188	1,270	1,442	1,524	1,727	1,756	1,792	1,877
Deaths	110	124	118	140	143	167	152	122	124
Rate	99.3	111.7	99.0	103.3	99.3	102.4	91.6	70.9	68.7
Glomerular Diseases									
Patients	257	231	251	258	275	288	291	270	272
Deaths	14	16	16	23	19	21	14	15	7
Rate	56.3	72.1	66.8	94.9	71.6	76.2	49.4	56.9	26.0
Diabetes									
Patients	288	304	378	413	492	550	542	590	674
Deaths	37	43	51	46	54	70	62	47	57
Rate	136.7	155.3	147.1	119.4	116.4	137.5	123.8	83.3	89.1
Hypertensive Nephrosclerosis									
Patients	224	264	293	356	385	431	499	525	547
Deaths	23	29	30	44	36	41	45	34	38
Rate	110.3	119.3	110.2	133.8	99.5	100.7	96.1	67.7	72.4
Polycystic Kidneys									
Patients	108	120	112	123	112	156	167	135	142
Deaths	6	5	9	3	8	9	9	6	6
Rate	57.5	42.8	85.6	24.6	75.4	59.6	55.0	45.7	43.4
Tubular and Interstitial Diseases									
Patients	70	45	42	66	60	96	85	74	82
Deaths	2	12	2	8	6	7	6	5	8
Rate	29.1	311.6	48.4	130.1	107.4	76.3	73.7	71.0	104.2
Renovascular & Other Vascular Diseases									
Patients	91	65	82	80	64	60	52	58	47
Deaths	12	6	4	9	7	6	6	7	1
Rate	141.4	96.3	50.7	120.8	117.9	106.9	124.6	130.6	21.7

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.7b (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

Primary Diagnosis	Year of Transplant								
	2000	2001	2002	2003	2004	2005	2006	2007	2008
Congenital, Rare Familial, & Metabolic Disorders									
Patients	8	17	14	15	11	12	10	13	12
Deaths	0	1	0	1	2	1	1	1	0
Rate	*	60.7	0.0	70.3	196.8	84.5	104.1	78.4	0.0
Neoplasms									
Patients	1	5	4	5	7	12	3	6	9
Deaths	0	0	0	1	1	1	0	0	1
Rate	*	*	*	*	*	90.4	*	*	*
Other									
Patients	100	116	82	119	104	113	97	114	91
Deaths	13	12	4	5	9	10	7	6	6
Rate	143.0	111.0	50.3	43.3	92.2	92.8	75.0	54.7	68.3
Unknown									
Patients	23	21	12	7	14	9	10	7	1
Deaths	3	0	2	0	1	1	2	1	0
Rate	139.0	0.0	186.3	*	74.8	*	233.2	*	*

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.7b (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

PRA at Transplant	Year of Transplant								
	2000	2001	2002	2003	2004	2005	2006	2007	2008
Total									
Patients	1,170	1,188	1,270	1,442	1,524	1,727	1,756	1,792	1,877
Deaths	110	124	118	140	143	167	152	122	124
Rate	99.3	111.7	99.0	103.3	99.3	102.4	91.6	70.9	68.7
0%-9%									
Patients	897	915	1,049	1,165	1,221	1,430	1,459	1,463	1,514
Deaths	92	97	95	113	108	140	126	104	107
Rate	108.8	113.7	96.4	103.1	93.2	103.9	91.4	74.2	73.7
10%-79%									
Patients	138	126	99	126	149	180	206	212	236
Deaths	9	9	14	15	18	14	21	9	12
Rate	68.1	74.2	157.0	129.2	131.9	81.2	108.3	43.6	52.5
80%+									
Patients	38	36	34	28	26	24	26	37	45
Deaths	3	1	3	1	1	2	1	6	3
Rate	81.2	28.1	92.0	35.8	39.8	88.0	39.8	176.6	69.6
Unknown									
Patients	97	111	88	123	128	93	65	80	82
Deaths	6	17	6	11	16	11	4	3	2
Rate	64.6	169.8	70.8	95.2	133.2	126.2	64.5	38.2	24.7

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.7b (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

Donor Age	Year of Transplant								
	2000	2001	2002	2003	2004	2005	2006	2007	2008
Total									
Patients	1,170	1,188	1,270	1,442	1,524	1,727	1,756	1,792	1,877
Deaths	110	124	118	140	143	167	152	122	124
Rate	99.3	111.7	99.0	103.3	99.3	102.4	91.6	70.9	68.7
50-64 Years									
Patients	913	881	948	1,097	1,171	1,356	1,322	1,380	1,449
Deaths	84	73	76	102	111	123	104	86	92
Rate	97.0	87.4	84.4	98.6	100.4	95.9	82.9	64.7	66.1
65+ Years									
Patients	257	307	322	345	353	371	434	412	428
Deaths	26	51	42	38	32	44	48	36	32
Rate	107.6	185.6	144.0	118.5	95.5	126.6	118.7	92.2	77.6

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.7b (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

Level of HLA Mismatch (Total)		Year of Transplant								
		2000	2001	2002	2003	2004	2005	2006	2007	2008
Total										
	Patients	1,170	1,188	1,270	1,442	1,524	1,727	1,756	1,792	1,877
	Deaths	110	124	118	140	143	167	152	122	124
	Rate	99.3	111.7	99.0	103.3	99.3	102.4	91.6	70.9	68.7
A Locus Mismatch										
	0									
	Patients	247	231	188	175	180	212	226	225	233
	Deaths	19	20	20	12	10	19	20	13	14
	Rate	79.9	91.8	115.9	71.8	57.8	94.6	93.8	59.6	62.8
	1									
	Patients	428	477	492	536	606	669	661	704	720
	Deaths	39	47	48	55	53	61	60	53	46
	Rate	96.7	104.2	103.9	109.9	91.8	96.4	96.0	78.8	66.2
	2									
	Patients	495	480	590	730	738	845	869	863	924
	Deaths	52	57	50	73	80	87	72	56	64
	Rate	111.5	129.2	89.7	106.3	115.9	109.3	87.6	67.5	72.3
	Unknown									
	Patients	0	0	0	1	0	1	0	0	0
	Deaths	0	0	0	0	0	0	0	0	0
	Rate	0.0	0.0	0.0	*	0.0	*	0.0	0.0	0.0

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.7b (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

Level of HLA Mismatch (Total)		Year of Transplant								
		2000	2001	2002	2003	2004	2005	2006	2007	2008
Total										
	Patients	1,170	1,188	1,270	1,442	1,524	1,727	1,756	1,792	1,877
	Deaths	110	124	118	140	143	167	152	122	124
	Rate	99.3	111.7	99.0	103.3	99.3	102.4	91.6	70.9	68.7
B Locus Mismatch										
	0									
	Patients	252	219	177	130	119	156	136	144	155
	Deaths	21	19	20	9	7	13	10	10	7
	Rate	86.7	91.2	122.5	72.5	60.8	88.2	77.3	72.0	46.5
	1									
	Patients	435	427	413	374	400	399	372	414	433
	Deaths	36	45	28	42	32	50	36	23	28
	Rate	86.7	112.9	70.6	120.0	83.4	136.6	103.4	57.4	67.1
	2									
	Patients	483	542	680	937	1,005	1,171	1,248	1,234	1,289
	Deaths	53	60	70	89	104	104	106	89	89
	Rate	117.8	119.3	110.7	101.1	110.5	93.2	89.7	75.4	72.0
	Unknown									
	Patients	-	-	-	1	-	1	-	-	-
	Deaths	-	-	-	0	-	0	-	-	-
	Rate	-	-	-	*	-	*	-	-	-

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.7b (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

Level of HLA Mismatch (Total)		Year of Transplant								
		2000	2001	2002	2003	2004	2005	2006	2007	2008
Total										
	Patients	1,170	1,188	1,270	1,442	1,524	1,727	1,756	1,792	1,877
	Deaths	110	124	118	140	143	167	152	122	124
	Rate	99.3	111.7	99.0	103.3	99.3	102.4	91.6	70.9	68.7
DR Locus Mismatch										
	0									
	Patients	322	307	299	227	169	216	206	213	217
	Deaths	23	15	31	22	12	21	13	14	11
	Rate	74.1	50.5	111.3	103.6	74.0	103.8	65.9	68.3	52.4
	1									
	Patients	526	512	519	570	636	697	713	707	722
	Deaths	57	67	40	53	63	52	67	53	53
	Rate	115.6	142.1	81.0	98.6	105.5	78.1	100.0	78.4	76.6
	2									
	Patients	321	369	452	644	719	813	837	872	938
	Deaths	30	42	47	64	68	94	72	55	60
	Rate	99.0	123.0	112.0	105.8	99.9	123.5	90.8	65.6	66.5
	Unknown									
	Patients	1	-	-	1	-	1	-	-	-
	Deaths	0	-	-	1	-	0	-	-	-
	Rate	*	-	-	*	-	*	-	-	-

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.7b (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

Level of HLA Mismatch (Total)		Year of Transplant								
		2000	2001	2002	2003	2004	2005	2006	2007	2008
Total										
	Patients	1,170	1,188	1,270	1,442	1,524	1,727	1,756	1,792	1,877
	Deaths	110	124	118	140	143	167	152	122	124
	Rate	99.3	111.7	99.0	103.3	99.3	102.4	91.6	70.9	68.7
Total Mismatch										
0										
	Patients	136	122	103	93	86	123	110	124	115
	Deaths	12	5	10	7	3	11	6	10	6
	Rate	92.2	42.0	104.0	79.6	35.6	95.4	56.7	84.2	53.9
1										
	Patients	61	36	26	21	5	8	4	7	12
	Deaths	4	4	6	2	0	0	1	0	0
	Rate	67.6	119.4	282.6	100.2	*	*	*	*	0.0
2										
	Patients	108	124	95	55	51	39	48	43	52
	Deaths	7	13	8	5	2	2	5	0	4
	Rate	66.5	111.9	88.9	95.7	40.3	52.9	109.3	0.0	80.9
3										
	Patients	224	242	228	164	162	182	160	160	177
	Deaths	16	26	12	15	15	18	21	9	9
	Rate	74.2	115.0	54.5	97.1	97.9	105.8	144.6	58.7	52.4
4										
	Patients	283	282	322	343	408	427	450	463	471
	Deaths	34	31	32	35	41	40	32	35	35
	Rate	130.4	117.2	105.7	108.5	106.4	99.0	74.5	78.8	77.7
5										
	Patients	238	232	296	482	531	599	630	632	654
	Deaths	22	25	36	49	53	61	61	46	42
	Rate	98.1	116.8	132.5	109.0	106.1	108.2	103.0	76.1	66.7

(Continued)

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDDs.

Table 5.7b (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

Level of HLA Mismatch (Total)	Year of Transplant								
	2000	2001	2002	2003	2004	2005	2006	2007	2008
Total Mismatch									
6									
Patients	119	150	200	282	281	348	354	363	396
Deaths	15	20	14	26	29	35	26	22	28
Rate	135.1	146.2	73.7	97.2	110.1	106.1	77.0	63.0	73.8
Unknown									
Patients	1	-	-	2	-	1	-	-	-
Deaths	0	-	-	1	-	0	-	-	-
Rate	*	-	-	*	-	*	-	-	-

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.7b (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

Cold Ischemic Time	Year of Transplant								
	2000	2001	2002	2003	2004	2005	2006	2007	2008
Total									
Patients	1,170	1,188	1,270	1,442	1,524	1,727	1,756	1,792	1,877
Deaths	110	124	118	140	143	167	152	122	124
Rate	99.3	111.7	99.0	103.3	99.3	102.4	91.6	70.9	68.7
0-11 Hours									
Patients	105	123	182	254	256	302	336	356	367
Deaths	7	15	13	22	20	25	20	21	19
Rate	69.0	133.0	75.6	91.9	82.1	87.2	61.8	60.9	53.7
12-21 Hours									
Patients	432	475	514	584	600	704	755	740	796
Deaths	38	43	44	50	46	72	73	53	42
Rate	92.2	95.8	90.4	90.8	80.0	108.3	102.9	75.0	54.4
22-31 Hours									
Patients	341	307	291	302	365	427	382	394	456
Deaths	30	33	35	27	38	37	37	29	45
Rate	92.9	115.4	130.5	94.2	110.9	91.4	102.4	76.8	105.0
32-41 Hours									
Patients	94	84	80	78	93	95	121	147	109
Deaths	13	8	7	9	6	5	10	7	4
Rate	149.2	100.6	92.6	123.4	67.1	53.9	88.2	48.8	37.8
42+ Hours									
Patients	19	14	16	32	27	32	37	51	52
Deaths	3	1	0	4	5	2	2	3	5
Rate	169.7	75.3	0.0	134.1	211.4	65.8	56.7	61.5	100.4
Unknown									
Patients	179	185	187	192	183	167	125	104	97
Deaths	19	24	19	28	28	26	10	9	9
Rate	114.4	141.4	109.4	159.1	169.0	173.0	85.6	91.7	96.3

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.7b (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

Transplant Center State	Year of Transplant								
	2000	2001	2002	2003	2004	2005	2006	2007	2008
Total									
Patients	1,170	1,188	1,270	1,442	1,524	1,727	1,756	1,792	1,877
Deaths	110	124	118	140	143	167	152	122	124
Rate	99.3	111.7	99.0	103.3	99.3	102.4	91.6	70.9	68.7
Alabama									
Patients	37	25	18	12	19	32	6	21	25
Deaths	0	1	1	1	5	3	0	3	2
Rate	0.0	40.1	57.8	85.6	329.7	101.6	*	157.1	83.1
Arkansas									
Patients	4	5	6	4	6	13	16	11	5
Deaths	1	0	1	0	0	2	1	1	0
Rate	*	*	*	*	*	167.1	65.6	97.2	*
Arizona									
Patients	11	11	13	12	22	17	17	30	44
Deaths	1	1	0	1	0	3	2	1	1
Rate	91.1	97.9	0.0	85.6	0.0	202.6	125.0	34.1	23.1
California									
Patients	148	150	132	208	212	160	172	179	193
Deaths	10	22	14	21	22	17	12	10	15
Rate	70.4	163.0	112.3	107.7	111.0	111.8	72.8	57.8	81.7
Colorado									
Patients	5	3	13	16	9	7	12	10	15
Deaths	0	0	0	1	0	1	1	0	0
Rate	*	*	0.0	66.2	*	*	84.5	0.0	0.0
Connecticut									
Patients	7	9	17	9	2	12	15	16	17
Deaths	2	1	4	1	0	1	1	3	1
Rate	*	*	280.6	*	*	87.1	69.1	209.3	61.0
District of Columbia									
Patients	10	9	10	3	12	18	26	20	13
Deaths	0	2	1	0	0	2	3	0	0
Rate	0.0	*	110.6	*	0.0	119.4	126.5	0.0	0.0

(Continued)

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.7b (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

Transplant Center State	Year of Transplant								
	2000	2001	2002	2003	2004	2005	2006	2007	2008
Delaware									
Patients	-	-	-	-	-	-	-	3	3
Deaths	-	-	-	-	-	-	-	0	0
Rate	-	-	-	-	-	-	-	*	*
Florida									
Patients	49	71	61	61	73	121	94	99	117
Deaths	5	5	4	6	6	11	8	8	5
Rate	110.0	72.8	67.7	102.6	85.0	97.1	91.1	84.5	43.5
Georgia									
Patients	29	34	36	24	31	34	34	20	36
Deaths	1	5	4	4	5	1	7	0	2
Rate	34.9	156.6	120.6	177.6	180.9	29.7	239.6	0.0	57.8
Hawaii									
Patients	11	7	14	7	1	6	18	10	7
Deaths	0	0	2	0	0	0	0	1	0
Rate	0.0	*	159.5	*	*	*	0.0	107.3	*
Iowa									
Patients	16	5	13	6	4	16	4	13	7
Deaths	2	1	0	0	0	2	0	2	2
Rate	133.7	*	0.0	*	*	135.0	*	167.8	*
Illinois									
Patients	41	43	38	92	98	62	82	64	82
Deaths	4	3	4	6	13	9	11	5	11
Rate	100.9	71.7	112.1	68.3	143.7	158.5	143.1	82.0	146.7
Indiana									
Patients	3	4	10	17	15	21	10	4	15
Deaths	0	0	0	0	1	0	1	0	1
Rate	*	*	0.0	0.0	67.2	0.0	104.0	*	67.5
Kansas									
Patients	-	1	1	5	3	3	2	11	10
Deaths	-	0	0	0	0	1	0	0	1
Rate	-	*	*	*	*	*	*	0.0	101.9

(Continued)

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.7b (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

Transplant Center State	Year of Transplant								
	2000	2001	2002	2003	2004	2005	2006	2007	2008
Kentucky									
Patients	12	15	20	19	9	2	15	19	9
Deaths	2	2	1	2	1	0	0	0	0
Rate	190.2	146.1	51.6	109.6	*	*	0.0	0.0	*
Louisiana									
Patients	10	19	16	13	23	29	27	28	33
Deaths	0	0	0	2	2	0	1	1	3
Rate	0.0	0.0	0.0	172.5	92.4	0.0	38.1	36.4	94.1
Massachusetts									
Patients	27	22	30	41	41	56	58	49	31
Deaths	3	1	1	6	3	5	1	3	1
Rate	118.7	46.3	34.0	157.9	75.3	94.3	17.5	64.1	32.8
Maryland									
Patients	109	83	66	65	58	51	54	77	63
Deaths	16	10	5	7	8	6	2	2	3
Rate	156.7	131.5	80.9	114.5	150.0	128.0	38.2	26.5	48.9
Maine									
Patients	8	11	2	1	3	3	1	1	2
Deaths	0	4	0	0	0	0	0	0	0
Rate	*	444.2	*	*	*	*	*	*	*
Michigan									
Patients	15	32	31	32	40	32	63	46	47
Deaths	1	2	2	5	5	4	7	4	3
Rate	67.2	65.7	67.8	172.6	132.7	135.2	121.3	90.5	66.5
Minnesota									
Patients	28	14	11	13	19	22	22	13	16
Deaths	3	0	1	3	2	2	0	1	2
Rate	115.7	0.0	98.3	271.6	108.3	99.3	0.0	80.0	134.3
Missouri									
Patients	16	7	6	13	9	24	27	30	12
Deaths	1	1	0	1	1	1	5	1	0
Rate	66.7	*	*	83.3	*	42.6	211.7	33.8	0.0

(Continued)

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.7b (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

Transplant Center State	Year of Transplant								
	2000	2001	2002	2003	2004	2005	2006	2007	2008
Mississippi									
Patients	-	1	1	-	-	-	-	2	14
Deaths	-	0	1	-	-	-	-	0	1
Rate	-	*	*	-	-	-	-	*	76.6
North Carolina									
Patients	31	23	45	28	48	59	45	73	80
Deaths	2	0	4	2	4	0	3	7	7
Rate	66.4	0.0	96.1	76.1	89.6	0.0	69.3	102.3	92.3
North Dakota									
Patients	5	9	12	6	5	7	5	2	5
Deaths	1	1	2	1	1	2	1	0	0
Rate	*	*	198.2	*	*	*	*	*	*
Nebraska									
Patients	6	10	4	7	8	10	10	15	12
Deaths	0	0	0	1	0	1	1	1	0
Rate	*	0.0	*	*	*	105.8	104.1	69.6	0.0
New Hampshire									
Patients	2	3	1	6	1	1	14	5	4
Deaths	0	1	0	3	0	0	1	0	0
Rate	*	*	*	*	*	*	74.9	*	*
New Jersey									
Patients	41	55	54	43	77	79	52	57	73
Deaths	6	4	3	4	11	11	7	9	9
Rate	160.3	77.6	58.7	97.3	159.5	152.4	148.2	175.8	133.1
New Mexico									
Patients	8	5	2	-	7	14	3	10	6
Deaths	0	0	0	-	0	1	0	1	0
Rate	*	*	*	-	*	75.8	*	104.2	*
Nevada									
Patients	2	5	4	7	-	-	2	2	3
Deaths	1	0	2	0	-	-	0	0	1
Rate	*	*	*	*	-	-	*	*	*

(Continued)

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.7b (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

Transplant Center State	Year of Transplant								
	2000	2001	2002	2003	2004	2005	2006	2007	2008
New York									
Patients	77	91	108	111	127	207	213	226	230
Deaths	9	10	16	14	12	24	15	14	11
Rate	124.3	118.3	161.2	138.7	99.6	124.6	74.0	64.5	49.2
Ohio									
Patients	51	32	55	58	46	73	76	57	74
Deaths	6	3	3	6	5	16	12	3	7
Rate	129.5	100.3	57.0	110.9	116.5	252.0	177.4	54.9	101.0
Oklahoma									
Patients	14	11	10	11	15	10	17	14	17
Deaths	1	2	1	2	0	2	2	1	0
Rate	72.4	217.1	101.7	211.1	0.0	221.9	129.5	75.6	0.0
Oregon									
Patients	21	20	33	22	30	11	19	22	14
Deaths	1	2	1	3	4	0	2	2	0
Rate	49.4	105.8	30.9	152.0	149.6	0.0	110.0	98.7	0.0
Pennsylvania									
Patients	108	116	157	182	122	158	204	157	157
Deaths	10	12	20	18	11	14	18	13	13
Rate	98.7	109.4	139.6	105.1	95.3	93.2	93.7	85.9	86.4
Puerto Rico									
Patients	2	5	6	-	5	15	6	17	16
Deaths	0	0	0	-	0	1	0	1	1
Rate	*	*	*	-	*	67.1	*	60.7	65.1
Rhode Island									
Patients	7	8	13	7	7	5	4	-	6
Deaths	2	1	1	0	1	0	0	-	1
Rate	*	*	78.6	*	*	*	*	-	*
South Carolina									
Patients	4	9	4	11	12	10	11	18	17
Deaths	1	4	0	3	2	1	0	1	1
Rate	*	*	*	351.7	198.3	100.6	0.0	58.4	59.8

(Continued)

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.7b (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

Transplant Center State	Year of Transplant								
	2000	2001	2002	2003	2004	2005	2006	2007	2008
South Dakota									
Patients	3	-	11	11	13	17	7	6	11
Deaths	0	-	1	1	0	0	2	0	1
Rate	*	-	99.8	97.6	0.0	0.0	*	*	95.5
Tennessee									
Patients	22	34	25	24	28	35	31	45	30
Deaths	2	4	2	2	3	1	1	7	2
Rate	94.1	127.4	83.3	86.2	115.0	28.6	33.3	172.3	71.4
Texas									
Patients	65	63	55	111	117	124	96	141	119
Deaths	4	5	3	4	3	12	8	7	6
Rate	65.1	84.9	56.4	37.0	25.9	102.7	88.2	51.4	52.1
Utah									
Patients	5	2	3	4	3	3	5	8	3
Deaths	0	0	0	0	0	0	1	1	0
Rate	*	*	*	*	*	*	*	*	*
Virginia									
Patients	15	26	35	35	46	47	68	53	74
Deaths	2	1	6	3	2	0	4	5	8
Rate	144.5	39.7	196.5	91.5	44.3	0.0	60.2	99.1	115.4
Vermont									
Patients	3	-	-	4	4	5	8	2	6
Deaths	1	-	-	1	0	0	1	0	0
Rate	*	-	-	*	*	*	*	*	*
Washington									
Patients	28	21	20	16	22	32	17	22	30
Deaths	4	1	3	0	2	5	0	1	0
Rate	152.0	49.3	167.6	0.0	93.5	169.8	0.0	45.6	0.0
Wisconsin									
Patients	45	54	46	62	65	62	64	63	69
Deaths	4	12	4	5	8	4	10	2	2
Rate	91.0	248.9	92.2	84.8	129.4	65.5	168.3	32.3	29.7

(Continued)

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.7b (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 2000 to 2008
Rates for Recipients during First Year after Deceased Donor ECD Kidney Transplantation

Transplant Center State	Year of Transplant								
	2000	2001	2002	2003	2004	2005	2006	2007	2008
West Virginia									
Patients	9	5	2	3	7	2	4	1	5
Deaths	1	0	0	0	0	1	0	0	0
Rate	*	*	*	*	*	*	*	*	*

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.