

**Table 5.12a**  
**Adjusted Patient Survival, Deceased Donor non-ECD Kidney Transplants**  
**Survival at 3 Months, 1 Year, 5 Years, and 10 Years**

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2007 - 2008)			(Tx 2007 - 2008)			(Tx 2003 - 2008)			(Tx 1998 - 2008)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
<b>Total</b>												
All	14,326	98.7%	0.1%	14,326	97.2%	0.2%	40,616	86.5%	0.2%	69,684	65.7%	0.3%
<b>Age at Tx</b>												
<1 Year	0	-	-	0	-	-	4	+	+	5	+	+
1-5 Years	134	98.0%	1.2%	134	97.3%	1.4%	355	94.1%	1.6%	551	91.4%	1.6%
6-11 Years	191	100.0%	0.0%	191	99.6%	0.4%	549	98.2%	0.6%	861	93.5%	1.4%
12-17 Years	520	99.7%	0.2%	520	99.1%	0.4%	1,484	95.0%	0.8%	2,179	85.2%	1.4%
18-34 Years	1,628	99.2%	0.2%	1,628	98.4%	0.3%	5,022	93.2%	0.5%	9,602	83.1%	0.6%
35-49 Years	4,091	99.1%	0.1%	4,091	98.3%	0.2%	11,708	90.0%	0.4%	21,306	72.8%	0.5%
50-64 Years	5,644	98.3%	0.2%	5,644	96.1%	0.3%	16,033	82.9%	0.4%	26,827	56.6%	0.5%
65+ Years	2,118	97.5%	0.3%	2,118	93.9%	0.5%	5,461	71.7%	0.8%	8,353	36.2%	1.0%
<b>Recipient Ethnicity/Race</b>												
White	6,317	98.9%	0.1%	6,317	97.3%	0.2%	18,377	85.9%	0.3%	33,420	64.6%	0.4%
African American	4,653	98.3%	0.2%	4,653	96.5%	0.3%	12,919	84.8%	0.4%	21,539	61.9%	0.6%
Hispanic/Latino	2,313	99.1%	0.2%	2,313	97.8%	0.3%	6,265	89.8%	0.5%	9,827	71.8%	0.8%
Asian	862	98.6%	0.4%	862	97.7%	0.5%	2,499	90.8%	0.7%	4,017	77.0%	1.1%
Other/Multi-race	181	99.2%	0.6%	181	98.3%	0.8%	556	87.5%	1.7%	879	71.5%	2.4%
Unknown	0	-	-	0	-	-	0	-	-	2	+	+
<b>Recipient Gender</b>												
Female	5,617	98.7%	0.1%	5,617	97.3%	0.2%	16,078	87.6%	0.3%	27,742	67.1%	0.5%
Male	8,709	98.8%	0.1%	8,709	97.1%	0.2%	24,538	85.7%	0.3%	41,942	64.7%	0.4%

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether kidneys were allocated under ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

**Table 5.12a (Continued)**  
**Adjusted Patient Survival, Deceased Donor non-ECD Kidney Transplants**  
**Survival at 3 Months, 1 Year, 5 Years, and 10 Years**

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2007 - 2008)			(Tx 2007 - 2008)			(Tx 2003 - 2008)			(Tx 1998 - 2008)		
	N	%	Std. Err.									
<b>Total</b>												
All	14,326	98.7%	0.1%	14,326	97.2%	0.2%	40,616	86.5%	0.2%	69,684	65.7%	0.3%
<b>Primary Diagnosis</b>												
Glomerular Diseases	3,276	99.0%	0.2%	3,276	97.9%	0.3%	9,593	89.4%	0.4%	17,255	73.9%	0.6%
Diabetes	3,580	98.5%	0.2%	3,580	95.7%	0.3%	9,814	80.2%	0.5%	16,303	50.0%	0.7%
Hypertensive												
Nephrosclerosis	3,647	98.5%	0.2%	3,647	97.0%	0.3%	9,607	85.0%	0.5%	15,047	63.5%	0.7%
Polycystic Kidneys	1,228	99.7%	0.2%	1,228	98.7%	0.3%	3,524	93.4%	0.5%	6,347	79.5%	0.8%
Tubular and Interstitial Diseases	607	98.8%	0.4%	607	97.7%	0.6%	1,946	86.8%	1.0%	3,581	68.4%	1.2%
Renovascular & Other Vascular Diseases	413	97.9%	0.7%	413	95.5%	1.0%	1,606	87.6%	0.9%	3,335	62.8%	1.3%
Congenital, Rare Familial, & Metabolic Disorders	422	99.0%	0.6%	422	97.6%	1.0%	1,167	90.6%	1.3%	1,898	76.7%	1.9%
Neoplasms	55	96.9%	2.1%	55	96.9%	2.1%	148	82.8%	4.0%	226	52.1%	5.9%
Other	1,091	98.2%	0.4%	1,091	97.4%	0.5%	3,192	88.6%	0.7%	5,664	69.0%	1.0%
Unknown	7	+	+	7	+	+	19	59.3%	15.3%	28	73.6%	9.3%

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether kidneys were allocated under ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

**Table 5.12a (Continued)**  
**Adjusted Patient Survival, Deceased Donor non-ECD Kidney Transplants**  
**Survival at 3 Months, 1 Year, 5 Years, and 10 Years**

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2007 - 2008)			(Tx 2007 - 2008)			(Tx 2003 - 2008)			(Tx 1998 - 2008)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
<b>Total</b>												
All	14,326	98.7%	0.1%	14,326	97.2%	0.2%	40,616	86.5%	0.2%	69,684	65.7%	0.3%
<b>Transplant Center State</b>												
Alabama	248	99.2%	0.6%	248	98.4%	0.8%	651	84.9%	1.9%	1,275	63.1%	2.0%
Arkansas	140	99.3%	0.6%	140	98.0%	1.1%	407	86.9%	2.2%	598	56.4%	4.1%
Arizona	255	99.7%	0.3%	255	98.2%	0.7%	641	90.9%	1.5%	1,048	73.1%	2.2%
California	1,689	99.2%	0.2%	1,689	98.0%	0.3%	4,938	87.9%	0.6%	8,224	69.4%	0.9%
Colorado	146	99.5%	0.5%	146	97.3%	1.2%	528	89.7%	1.6%	885	66.4%	2.8%
Connecticut	108	99.2%	0.8%	108	99.2%	0.8%	336	89.5%	2.0%	564	67.7%	3.3%
District of Columbia	131	99.3%	0.7%	131	98.6%	0.9%	433	86.1%	2.1%	708	67.4%	2.9%
Delaware	17	+	+	17	+	+	24	+	+	30	+	+
Florida	1,036	98.9%	0.3%	1,036	98.0%	0.4%	2,789	86.4%	0.8%	4,772	65.4%	1.1%
Georgia	362	98.1%	0.7%	362	96.3%	0.9%	1,075	86.3%	1.3%	1,914	65.9%	1.8%
Hawaii	48	95.9%	2.3%	48	94.6%	2.6%	149	91.3%	2.5%	253	76.0%	4.5%
Iowa	120	98.8%	0.8%	120	96.3%	1.5%	324	86.9%	2.2%	594	61.9%	3.1%
Illinois	449	98.6%	0.5%	449	97.6%	0.7%	1,378	86.8%	1.1%	2,510	64.7%	1.6%
Indiana	198	98.4%	0.8%	198	98.1%	0.9%	679	88.9%	1.4%	1,188	76.4%	1.9%
Kansas	140	98.8%	0.9%	140	96.3%	1.5%	362	90.9%	1.7%	602	71.9%	3.0%
Kentucky	134	100.0%	0.0%	134	99.3%	0.6%	447	84.8%	2.1%	829	59.6%	2.7%
Louisiana	264	98.6%	0.7%	264	96.9%	1.0%	865	83.9%	1.5%	1,441	59.8%	2.4%
Massachusetts	255	96.5%	1.1%	255	95.5%	1.2%	773	89.0%	1.3%	1,405	67.1%	2.0%
Maryland	282	98.2%	0.7%	282	96.4%	1.0%	762	84.0%	1.6%	1,599	57.8%	1.9%
Maine	43	100.0%	0.0%	43	96.0%	2.8%	154	86.2%	3.1%	270	66.8%	4.4%
Michigan	525	97.8%	0.6%	525	95.9%	0.8%	1,388	83.6%	1.3%	2,357	63.0%	1.6%
Minnesota	181	100.0%	0.0%	181	99.1%	0.6%	583	88.7%	1.4%	1,046	70.2%	2.0%
Missouri	419	100.0%	0.0%	419	98.4%	0.6%	1,124	87.7%	1.3%	1,938	67.1%	1.8%
Mississippi	156	97.4%	1.3%	156	97.4%	1.3%	233	88.1%	3.2%	364	73.7%	4.4%
North Carolina	511	98.4%	0.5%	511	97.2%	0.7%	1,272	85.8%	1.4%	1,944	67.4%	1.9%
North Dakota	33	97.6%	2.4%	33	95.1%	3.3%	93	79.8%	4.6%	170	55.4%	6.4%
Nebraska	65	100.0%	0.0%	65	96.6%	1.9%	211	84.5%	2.9%	371	70.8%	3.6%
New Hampshire	28	100.0%	0.0%	28	97.3%	2.6%	88	88.2%	3.4%	139	82.5%	3.9%
New Jersey	335	96.0%	1.0%	335	93.4%	1.3%	845	82.3%	1.6%	1,429	60.5%	2.2%
New Mexico	82	99.0%	1.0%	82	98.0%	1.4%	223	95.3%	1.7%	363	74.2%	4.0%
Nevada	78	100.0%	0.0%	78	95.3%	2.3%	245	80.1%	3.1%	418	56.0%	3.8%

(Continued)

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether kidneys were allocated under ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

**Table 5.12a (Continued)**  
**Adjusted Patient Survival, Deceased Donor non-ECD Kidney Transplants**  
**Survival at 3 Months, 1 Year, 5 Years, and 10 Years**

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2007 - 2008)			(Tx 2007 - 2008)			(Tx 2003 - 2008)			(Tx 1998 - 2008)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
<b>Transplant Center State</b>												
New York	876	98.7%	0.3%	876	96.0%	0.6%	2,315	85.2%	1.0%	3,824	65.8%	1.3%
Ohio	527	98.9%	0.4%	527	96.7%	0.7%	1,388	83.1%	1.2%	2,661	60.4%	1.5%
Oklahoma	135	100.0%	0.0%	135	97.3%	1.3%	424	82.2%	2.2%	776	63.1%	2.6%
Oregon	155	99.5%	0.5%	155	97.8%	1.1%	489	88.0%	1.9%	828	69.8%	2.6%
Pennsylvania	938	98.6%	0.3%	938	96.9%	0.5%	2,826	84.5%	0.8%	4,811	64.5%	1.1%
Puerto Rico	116	99.2%	0.8%	116	99.2%	0.8%	339	90.8%	2.0%	511	62.7%	4.5%
Rhode Island	25	100.0%	0.0%	25	95.9%	3.9%	123	84.7%	3.5%	241	50.9%	5.7%
South Carolina	210	100.0%	0.0%	210	97.5%	1.0%	582	88.4%	1.7%	1,013	62.0%	2.6%
South Dakota	33	100.0%	0.0%	33	97.7%	2.3%	92	91.9%	3.2%	178	78.3%	3.7%
Tennessee	391	98.7%	0.5%	391	96.5%	0.9%	1,134	85.3%	1.3%	1,899	58.3%	2.0%
Texas	1,229	98.4%	0.3%	1,229	96.9%	0.5%	3,502	87.2%	0.7%	6,017	67.3%	1.0%
Utah	105	100.0%	0.0%	105	99.2%	0.8%	310	93.1%	1.7%	553	75.4%	2.8%
Virginia	340	97.3%	0.8%	340	95.8%	1.0%	894	85.1%	1.6%	1,488	64.4%	2.0%
Vermont	30	100.0%	0.0%	30	94.8%	3.5%	85	83.9%	5.2%	116	78.4%	5.7%
Washington	314	99.4%	0.4%	314	98.9%	0.6%	916	90.7%	1.1%	1,495	66.5%	2.2%
Wisconsin	362	98.4%	0.6%	362	96.3%	0.9%	991	86.7%	1.3%	1,662	66.0%	1.9%
West Virginia	62	97.2%	1.9%	62	95.8%	2.4%	186	77.0%	3.8%	363	54.5%	3.9%

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether kidneys were allocated under ECD system. Non-ECDs include all other donors, including SCDs and DCDs.