

**Table 5.12c**  
**Adjusted Patient Survival, Deceased Donor Kidney Transplants**  
**Survival at 3 Months, 1 Year, 5 Years, and 10 Years**

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2007 - 2008)			(Tx 2007 - 2008)			(Tx 2003 - 2008)			(Tx 1998 - 2008)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
<b>Total</b>												
All	17,760	98.6%	0.1%	17,760	96.6%	0.1%	49,989	84.2%	0.2%	84,554	61.8%	0.3%
<b>Age at Tx</b>												
<1 Year	0	-	-	0	-	-	4	+	+	5	+	+
1-5 Years	134	98.0%	1.2%	134	97.3%	1.4%	355	94.1%	1.6%	552	91.4%	1.6%
6-11 Years	191	100.0%	0.0%	191	99.6%	0.4%	551	98.2%	0.6%	868	93.4%	1.4%
12-17 Years	520	99.7%	0.2%	520	99.1%	0.4%	1,487	95.0%	0.8%	2,202	85.3%	1.4%
18-34 Years	1,679	99.2%	0.2%	1,679	98.4%	0.3%	5,248	92.9%	0.5%	10,222	82.2%	0.6%
35-49 Years	4,439	99.1%	0.1%	4,439	98.1%	0.2%	12,917	89.4%	0.4%	23,800	71.5%	0.5%
50-64 Years	7,417	98.3%	0.1%	7,417	95.7%	0.2%	20,688	81.2%	0.4%	34,156	54.3%	0.5%
65+ Years	3,380	97.2%	0.3%	3,380	93.0%	0.4%	8,739	68.6%	0.7%	12,749	33.3%	0.9%
<b>Recipient Ethnicity/Race</b>												
White	7,874	98.7%	0.1%	7,874	96.5%	0.2%	22,800	83.4%	0.3%	40,672	60.4%	0.4%
African American	5,758	98.3%	0.2%	5,758	96.2%	0.3%	15,895	82.7%	0.4%	26,194	58.3%	0.6%
Hispanic/Latino	2,786	98.9%	0.2%	2,786	97.4%	0.3%	7,415	87.8%	0.5%	11,549	68.3%	0.8%
Asian	1,109	98.5%	0.3%	1,109	97.3%	0.4%	3,171	88.5%	0.7%	5,028	73.3%	1.1%
Other/Multi-race	233	99.0%	0.5%	233	98.1%	0.8%	708	85.6%	1.6%	1,109	66.6%	2.3%
Unknown	0	-	-	0	-	-	0	-	-	2	+	+
<b>Recipient Gender</b>												
Female	6,865	98.5%	0.1%	6,865	96.9%	0.2%	19,489	85.5%	0.3%	33,310	63.5%	0.5%
Male	10,895	98.6%	0.1%	10,895	96.4%	0.2%	30,500	83.4%	0.3%	51,244	60.7%	0.4%

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether kidneys were allocated under ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

**Table 5.12c (Continued)**  
**Adjusted Patient Survival, Deceased Donor Kidney Transplants**  
**Survival at 3 Months, 1 Year, 5 Years, and 10 Years**

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2007 - 2008)			(Tx 2007 - 2008)			(Tx 2003 - 2008)			(Tx 1998 - 2008)		
	N	%	Std. Err.									
<b>Total</b>												
All	17,760	98.6%	0.1%	17,760	96.6%	0.1%	49,989	84.2%	0.2%	84,554	61.8%	0.3%
<b>Primary Diagnosis</b>												
Glomerular Diseases	3,733	99.0%	0.2%	3,733	97.7%	0.3%	10,995	88.3%	0.4%	19,771	70.9%	0.5%
Diabetes	4,812	98.3%	0.2%	4,812	95.2%	0.3%	12,977	77.5%	0.5%	21,012	46.3%	0.6%
Hypertensive												
Nephrosclerosis	4,673	98.2%	0.2%	4,673	96.4%	0.3%	12,212	82.7%	0.5%	18,812	59.9%	0.7%
Polycystic Kidneys	1,482	99.4%	0.2%	1,482	98.3%	0.3%	4,298	92.1%	0.5%	7,662	77.1%	0.7%
Tubular and Interstitial												
Diseases	745	98.5%	0.4%	745	96.9%	0.6%	2,356	84.2%	0.9%	4,244	65.0%	1.1%
Renovascular & Other												
Vascular Diseases	514	97.8%	0.6%	514	95.1%	0.9%	1,947	85.4%	0.9%	4,015	59.3%	1.2%
Congenital, Rare												
Familial, & Metabolic												
Disorders	442	99.0%	0.6%	442	97.3%	1.0%	1,224	89.4%	1.4%	2,015	74.0%	2.0%
Neoplasms	70	97.7%	1.6%	70	96.6%	1.9%	189	79.2%	3.7%	284	52.2%	5.2%
Other	1,281	98.2%	0.4%	1,281	96.9%	0.5%	3,766	86.5%	0.7%	6,703	64.8%	1.0%
Unknown	8	+	+	8	+	+	25	60.6%	15.6%	36	75.9%	8.3%

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether kidneys were allocated under ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

**Table 5.12c (Continued)**  
**Adjusted Patient Survival, Deceased Donor Kidney Transplants**  
**Survival at 3 Months, 1 Year, 5 Years, and 10 Years**

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2007 - 2008)			(Tx 2007 - 2008)			(Tx 2003 - 2008)			(Tx 1998 - 2008)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
<b>Total</b>												
All	17,760	98.6%	0.1%	17,760	96.6%	0.1%	49,989	84.2%	0.2%	84,554	61.8%	0.3%
<b>Transplant Center State</b>												
Alabama	293	98.3%	0.7%	293	97.0%	1.0%	762	81.6%	1.9%	1,506	58.7%	1.9%
Arkansas	153	99.4%	0.6%	153	97.6%	1.2%	455	83.7%	2.4%	663	52.8%	4.1%
Arizona	325	99.8%	0.2%	325	98.1%	0.7%	777	89.8%	1.4%	1,234	71.3%	2.2%
California	2,047	99.0%	0.2%	2,047	97.3%	0.3%	6,016	85.6%	0.6%	9,947	64.8%	0.8%
Colorado	169	99.5%	0.5%	169	97.7%	1.0%	594	88.0%	1.6%	984	61.9%	2.8%
Connecticut	140	98.9%	0.8%	140	97.7%	1.1%	402	86.8%	2.0%	670	65.0%	3.2%
District of Columbia	164	99.5%	0.5%	164	98.9%	0.7%	525	85.8%	1.9%	848	62.7%	2.8%
Delaware	22	+	+	22	+	+	29	+	+	35	+	+
Florida	1,241	98.8%	0.3%	1,241	97.4%	0.4%	3,319	84.2%	0.8%	5,593	62.0%	1.1%
Georgia	417	98.1%	0.6%	417	96.3%	0.9%	1,247	84.2%	1.3%	2,224	62.8%	1.7%
Hawaii	63	96.7%	1.9%	63	94.4%	2.4%	196	89.7%	2.4%	350	72.6%	3.7%
Iowa	138	99.0%	0.7%	138	95.3%	1.5%	369	86.0%	2.1%	679	58.8%	3.0%
Illinois	585	98.0%	0.5%	585	96.0%	0.8%	1,818	83.4%	1.1%	3,157	60.9%	1.5%
Indiana	216	98.6%	0.7%	216	97.9%	0.9%	755	88.3%	1.3%	1,298	74.7%	1.9%
Kansas	160	98.9%	0.7%	160	96.3%	1.4%	394	89.1%	1.9%	637	70.1%	3.1%
Kentucky	162	100.0%	0.0%	162	99.5%	0.5%	518	84.1%	2.0%	961	57.6%	2.6%
Louisiana	321	98.5%	0.7%	321	96.2%	1.0%	1,009	82.5%	1.5%	1,643	57.0%	2.4%
Massachusetts	331	97.0%	0.9%	331	95.7%	1.0%	1,028	85.6%	1.3%	1,769	63.7%	1.9%
Maryland	409	98.4%	0.6%	409	96.5%	0.8%	1,088	80.5%	1.5%	2,278	51.9%	1.6%
Maine	46	100.0%	0.0%	46	96.4%	2.5%	164	86.3%	3.0%	304	63.8%	4.3%
Michigan	617	97.9%	0.5%	617	95.7%	0.7%	1,636	80.7%	1.2%	2,721	60.0%	1.6%
Minnesota	206	100.0%	0.0%	206	98.4%	0.8%	673	88.8%	1.3%	1,233	66.3%	2.0%
Missouri	458	100.0%	0.0%	458	98.4%	0.5%	1,233	86.6%	1.3%	2,113	64.2%	1.8%
Mississippi	172	96.7%	1.4%	172	96.7%	1.4%	249	87.2%	3.4%	389	68.6%	4.8%
North Carolina	655	97.7%	0.5%	655	96.3%	0.7%	1,586	83.8%	1.3%	2,404	63.2%	1.9%
North Dakota	40	98.1%	1.9%	40	96.1%	2.7%	122	77.1%	4.3%	227	51.0%	5.8%
Nebraska	88	100.0%	0.0%	88	96.7%	1.6%	264	83.3%	2.8%	449	66.5%	3.6%
New Hampshire	36	100.0%	0.0%	36	98.1%	1.8%	113	86.7%	3.2%	171	79.5%	3.9%
New Jersey	461	96.1%	0.8%	461	92.2%	1.2%	1,200	78.6%	1.4%	2,004	56.9%	1.9%
New Mexico	98	99.2%	0.8%	98	97.5%	1.4%	262	94.1%	1.8%	424	70.9%	4.0%
Nevada	83	98.8%	1.2%	83	94.1%	2.6%	258	78.9%	3.2%	446	52.9%	3.8%

(Continued)

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether kidneys were allocated under ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

**Table 5.12c (Continued)**  
**Adjusted Patient Survival, Deceased Donor Kidney Transplants**  
**Survival at 3 Months, 1 Year, 5 Years, and 10 Years**

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2007 - 2008)			(Tx 2007 - 2008)			(Tx 2003 - 2008)			(Tx 1998 - 2008)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
<b>Transplant Center State</b>												
New York	1,290	98.7%	0.3%	1,290	95.9%	0.5%	3,324	83.1%	0.9%	5,288	61.4%	1.2%
Ohio	646	98.9%	0.4%	646	96.4%	0.7%	1,739	80.7%	1.1%	3,241	56.4%	1.4%
Oklahoma	163	99.5%	0.5%	163	97.4%	1.2%	502	80.9%	2.1%	901	60.0%	2.5%
Oregon	187	98.7%	0.7%	187	97.4%	1.0%	594	85.6%	1.8%	1,028	66.3%	2.4%
Pennsylvania	1,222	98.7%	0.3%	1,222	96.2%	0.5%	3,700	81.6%	0.8%	6,238	59.9%	1.0%
Puerto Rico	149	99.4%	0.6%	149	98.1%	1.1%	398	88.8%	2.1%	588	59.6%	4.6%
Rhode Island	31	100.0%	0.0%	31	94.2%	3.9%	150	82.3%	3.4%	301	45.1%	5.2%
South Carolina	244	99.6%	0.4%	244	97.1%	1.0%	657	85.1%	1.8%	1,118	58.2%	2.6%
South Dakota	49	100.0%	0.0%	49	97.1%	2.0%	156	79.1%	3.7%	264	64.4%	4.4%
Tennessee	461	97.8%	0.6%	461	95.3%	0.9%	1,313	84.1%	1.3%	2,200	55.4%	1.9%
Texas	1,475	98.3%	0.3%	1,475	96.6%	0.5%	4,162	85.7%	0.7%	6,982	64.1%	1.0%
Utah	116	100.0%	0.0%	116	98.6%	1.0%	335	91.7%	1.8%	598	72.8%	2.9%
Virginia	459	97.2%	0.7%	459	94.4%	1.0%	1,191	82.2%	1.5%	1,905	58.5%	2.0%
Vermont	36	100.0%	0.0%	36	95.8%	2.9%	111	84.0%	4.4%	147	75.4%	5.7%
Washington	365	99.5%	0.3%	365	98.8%	0.5%	1,051	89.2%	1.1%	1,760	64.6%	2.0%
Wisconsin	483	98.5%	0.5%	483	96.7%	0.7%	1,338	83.0%	1.3%	2,218	61.3%	1.7%
West Virginia	68	97.5%	1.7%	68	96.2%	2.1%	207	78.1%	3.5%	416	52.5%	3.7%

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether kidneys were allocated under ECD system. Non-ECDs include all other donors, including SCDs and DCDs.