

Table 5.14a
Unadjusted Patient Survival, Deceased Donor non-ECD Kidney Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2007 - 2008)			(Tx 2007 - 2008)			(Tx 2003 - 2008)			(Tx 1998 - 2008)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	14,326	98.5%	0.1%	14,326	96.6%	0.2%	40,616	85.0%	0.2%	69,684	64.9%	0.3%
Age at Tx												
<1 Year	0	-	-	0	-	-	4	100.0%	0.0%	5	100.0%	0.0%
1-5 Years	134	97.8%	1.3%	134	97.0%	1.5%	355	94.1%	1.6%	551	91.4%	1.6%
6-11 Years	191	100.0%	0.0%	191	99.5%	0.5%	549	98.1%	0.6%	861	93.2%	1.5%
12-17 Years	520	99.6%	0.3%	520	98.8%	0.5%	1,484	94.9%	0.8%	2,179	84.1%	1.4%
18-34 Years	1,628	99.3%	0.2%	1,628	98.5%	0.3%	5,022	93.8%	0.4%	9,602	84.6%	0.5%
35-49 Years	4,091	99.1%	0.1%	4,091	98.3%	0.2%	11,708	90.1%	0.3%	21,306	73.2%	0.5%
50-64 Years	5,644	98.1%	0.2%	5,644	95.8%	0.3%	16,033	81.8%	0.4%	26,827	54.8%	0.5%
65+ Years	2,118	97.3%	0.4%	2,118	93.5%	0.5%	5,461	69.8%	0.8%	8,353	34.5%	1.0%
Recipient Ethnicity/Race												
White	6,317	98.6%	0.1%	6,317	96.6%	0.2%	18,377	83.6%	0.3%	33,420	62.7%	0.4%
African American	4,653	98.2%	0.2%	4,653	96.2%	0.3%	12,919	84.3%	0.4%	21,539	63.3%	0.5%
Hispanic/Latino	2,313	98.9%	0.2%	2,313	97.4%	0.3%	6,265	88.8%	0.5%	9,827	71.7%	0.8%
Asian	862	98.4%	0.4%	862	97.3%	0.5%	2,499	90.2%	0.7%	4,017	77.6%	1.1%
Other/Multi-race	181	98.9%	0.8%	181	97.8%	1.1%	556	84.9%	2.1%	879	63.6%	2.7%
Unknown	0	-	-	0	-	-	0	-	-	2	100.0%	0.0%
Recipient Gender												
Female	5,617	98.5%	0.2%	5,617	96.9%	0.2%	16,078	86.5%	0.3%	27,742	66.8%	0.5%
Male	8,709	98.5%	0.1%	8,709	96.5%	0.2%	24,538	84.1%	0.3%	41,942	63.6%	0.4%
Recipient Blood Type												
O	6,367	98.7%	0.1%	6,367	96.9%	0.2%	18,181	85.5%	0.3%	30,916	65.7%	0.4%
A	5,152	98.6%	0.2%	5,152	96.8%	0.2%	14,738	84.7%	0.4%	25,825	64.2%	0.5%
B	2,013	98.0%	0.3%	2,013	95.9%	0.4%	5,532	84.3%	0.6%	9,172	64.1%	0.8%
AB	794	97.6%	0.5%	794	95.5%	0.7%	2,165	85.3%	1.0%	3,771	65.1%	1.2%

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Dialysis needed within first week excludes patients whose graft failed before the end of the week.

Table 5.14a (Continued)
Unadjusted Patient Survival, Deceased Donor non-ECD Kidney Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2007 - 2008)			(Tx 2007 - 2008)			(Tx 2003 - 2008)			(Tx 1998 - 2008)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	14,326	98.5%	0.1%	14,326	96.6%	0.2%	40,616	85.0%	0.2%	69,684	64.9%	0.3%
Recipient Residency												
U.S. Resident	14,194	98.5%	0.1%	14,194	96.6%	0.2%	40,322	85.0%	0.2%	69,141	64.7%	0.3%
Nonresident Alien	132	98.5%	1.1%	132	97.0%	1.5%	288	95.8%	1.5%	494	91.3%	1.6%
Unknown	0	-	-	0	-	-	6	55.6%	24.8%	49	65.6%	7.1%
Hospitalized at Tx												
Not Hospitalized	14,156	98.6%	0.1%	14,156	96.7%	0.1%	40,127	85.1%	0.2%	68,591	64.9%	0.3%
Hospitalized	143	95.1%	1.8%	143	92.3%	2.2%	418	77.8%	2.5%	935	60.2%	2.2%
In Intensive Care	22	86.4%	7.3%	22	86.4%	7.3%	64	79.4%	5.1%	139	56.6%	5.2%
Unknown	5	100.0%	0.0%	5	100.0%	0.0%	7	85.7%	13.2%	19	80.2%	10.4%
Primary Diagnosis												
Glomerular Diseases	3,276	99.1%	0.2%	3,276	98.1%	0.2%	9,593	90.3%	0.4%	17,255	76.3%	0.5%
Diabetes	3,580	98.0%	0.2%	3,580	94.4%	0.4%	9,814	75.6%	0.6%	16,303	44.0%	0.7%
Hypertensive												
Nephrosclerosis	3,647	98.2%	0.2%	3,647	96.6%	0.3%	9,607	83.9%	0.5%	15,047	62.7%	0.7%
Polycystic Kidneys	1,228	99.6%	0.2%	1,228	98.5%	0.4%	3,524	91.8%	0.6%	6,347	75.8%	0.8%
Tubular and Interstitial Diseases	607	98.7%	0.5%	607	97.5%	0.6%	1,946	86.0%	1.0%	3,581	68.2%	1.1%
Renovascular & Other Vascular Diseases	413	97.6%	0.8%	413	94.9%	1.1%	1,606	86.8%	1.0%	3,335	63.0%	1.2%
Congenital, Rare Familial, & Metabolic Disorders	422	99.3%	0.4%	422	98.3%	0.6%	1,167	94.0%	0.8%	1,898	84.8%	1.3%
Neoplasms	55	96.4%	2.5%	55	96.4%	2.5%	148	79.7%	4.7%	226	46.3%	6.2%
Other	1,091	98.3%	0.4%	1,091	97.4%	0.5%	3,192	89.0%	0.7%	5,664	71.0%	0.9%
Unknown	7	100.0%	0.0%	7	100.0%	0.0%	19	65.7%	13.5%	28	75.8%	8.8%

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Dialysis needed within first week excludes patients whose graft failed before the end of the week.

Table 5.14a (Continued)
Unadjusted Patient Survival, Deceased Donor non-ECD Kidney Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2007 - 2008)			(Tx 2007 - 2008)			(Tx 2003 - 2008)			(Tx 1998 - 2008)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	14,326	98.5%	0.1%	14,326	96.6%	0.2%	40,616	85.0%	0.2%	69,684	64.9%	0.3%
Cold Ischemic Time												
0-11 Hours	3,254	98.5%	0.2%	3,254	96.6%	0.3%	8,539	86.2%	0.5%	12,761	66.9%	0.8%
12-21 Hours	6,088	98.6%	0.2%	6,088	97.0%	0.2%	16,896	85.3%	0.4%	28,621	64.9%	0.5%
22-31 Hours	3,233	98.5%	0.2%	3,233	96.1%	0.3%	9,189	84.7%	0.5%	16,690	64.2%	0.6%
32-41 Hours	612	98.2%	0.5%	612	95.4%	0.8%	1,680	83.4%	1.2%	3,353	60.7%	1.2%
42+ Hours	306	98.7%	0.6%	306	97.1%	1.0%	560	81.6%	2.6%	850	59.9%	2.7%
Unknown	833	98.1%	0.5%	833	97.0%	0.6%	3,752	83.4%	0.7%	7,409	65.4%	0.8%
PRA at Transplant												
0-9%	11,285	98.5%	0.1%	11,285	96.7%	0.2%	32,519	85.1%	0.3%	56,423	65.2%	0.3%
10-79%	1,864	98.4%	0.3%	1,864	96.3%	0.4%	4,689	84.0%	0.7%	7,046	61.7%	1.0%
80%+	675	97.8%	0.6%	675	95.7%	0.8%	1,681	85.3%	1.1%	2,342	61.5%	2.0%
Unknown	502	99.2%	0.4%	502	97.6%	0.7%	1,727	85.6%	1.0%	3,873	66.6%	1.1%
Yearly Center Volume												
0-12	276	99.6%	0.4%	276	97.8%	0.9%	932	90.0%	1.3%	1,872	69.9%	1.7%
13-28	788	99.0%	0.4%	788	96.3%	0.7%	2,515	85.5%	0.9%	4,847	66.5%	1.1%
29-47	1,708	98.0%	0.3%	1,708	96.0%	0.5%	4,655	83.2%	0.7%	9,038	63.2%	0.8%
48-81	2,263	98.4%	0.3%	2,263	96.1%	0.4%	8,403	85.2%	0.5%	15,539	63.5%	0.6%
82+	9,291	98.5%	0.1%	9,291	96.9%	0.2%	24,111	85.1%	0.3%	38,388	65.3%	0.4%
Dialysis Needed Within First Week After Tx												
No	10,980	98.8%	0.1%	10,980	97.4%	0.2%	31,410	87.0%	0.2%	54,072	67.4%	0.3%
Yes	3,338	97.5%	0.3%	3,338	94.2%	0.4%	9,195	78.4%	0.5%	15,516	55.8%	0.6%
Unknown	8	100.0%	0.0%	8	100.0%	0.0%	11	71.6%	14.0%	96	56.9%	5.3%
Donor Age												
<1 Year	98	98.0%	1.4%	98	96.9%	1.7%	185	84.5%	3.7%	326	63.2%	4.3%
1-5 Years	417	98.1%	0.7%	417	96.6%	0.9%	1,202	87.6%	1.1%	2,181	71.6%	1.4%
6-11 Years	419	98.6%	0.6%	419	96.9%	0.8%	1,209	86.5%	1.2%	2,517	66.7%	1.4%
12-17 Years	1,093	98.9%	0.3%	1,093	97.0%	0.5%	3,609	87.8%	0.7%	6,805	68.7%	0.8%
18-34 Years	5,159	98.9%	0.1%	5,159	97.5%	0.2%	14,422	87.5%	0.4%	24,128	68.0%	0.5%
35-49 Years	5,090	98.0%	0.2%	5,090	96.0%	0.3%	14,487	82.9%	0.4%	24,433	62.2%	0.5%
50-64 Years	2,050	98.5%	0.3%	2,050	95.8%	0.4%	5,502	81.3%	0.7%	9,294	58.4%	0.8%

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Dialysis needed within first week excludes patients whose graft failed before the end of the week.

Table 5.14a (Continued)
Unadjusted Patient Survival, Deceased Donor non-ECD Kidney Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2007 - 2008)			(Tx 2007 - 2008)			(Tx 2003 - 2008)			(Tx 1998 - 2008)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	14,326	98.5%	0.1%	14,326	96.6%	0.2%	40,616	85.0%	0.2%	69,684	64.9%	0.3%
A Locus Mismatch												
0	2,987	98.5%	0.2%	2,987	96.8%	0.3%	8,307	85.7%	0.5%	14,628	65.6%	0.6%
1	4,817	98.5%	0.2%	4,817	96.4%	0.3%	13,779	85.2%	0.4%	24,266	65.0%	0.5%
2	6,521	98.5%	0.1%	6,521	96.8%	0.2%	18,525	84.6%	0.3%	30,760	64.4%	0.4%
Unknown	1	100.0%	0.0%	1	100.0%	0.0%	5	80.0%	17.9%	30	52.6%	9.7%
B Locus Mismatch												
0	2,481	98.5%	0.2%	2,481	96.8%	0.4%	7,032	85.7%	0.5%	13,528	65.1%	0.6%
1	3,159	98.5%	0.2%	3,159	96.5%	0.3%	9,301	84.7%	0.5%	19,945	65.1%	0.5%
2	8,685	98.5%	0.1%	8,685	96.7%	0.2%	24,278	84.9%	0.3%	36,181	64.5%	0.4%
Unknown	1	100.0%	0.0%	1	100.0%	0.0%	5	80.0%	17.9%	30	52.6%	9.7%
DR Locus Mismatch												
0	3,743	98.6%	0.2%	3,743	96.8%	0.3%	10,749	85.4%	0.4%	19,931	64.9%	0.5%
1	6,216	98.4%	0.2%	6,216	96.7%	0.2%	17,561	85.0%	0.3%	29,626	65.5%	0.4%
2	4,366	98.5%	0.2%	4,366	96.5%	0.3%	12,288	84.7%	0.4%	20,066	64.0%	0.6%
Unknown	1	100.0%	0.0%	1	100.0%	0.0%	18	83.3%	8.8%	61	58.5%	7.3%
Total Mismatch												
0	2,217	98.4%	0.3%	2,217	96.8%	0.4%	6,167	85.6%	0.6%	10,295	65.5%	0.8%
1	114	99.1%	0.9%	114	97.4%	1.5%	364	86.4%	2.2%	1,399	63.9%	1.6%
2	538	98.5%	0.5%	538	96.1%	0.8%	1,565	83.9%	1.1%	4,313	64.8%	1.0%
3	1,740	98.5%	0.3%	1,740	96.4%	0.4%	5,069	85.7%	0.6%	10,869	65.8%	0.6%
4	3,552	98.8%	0.2%	3,552	96.9%	0.3%	10,302	85.0%	0.5%	16,852	64.6%	0.6%
5	4,266	98.2%	0.2%	4,266	96.5%	0.3%	11,900	85.0%	0.4%	17,622	64.6%	0.6%
6	1,898	98.6%	0.3%	1,898	96.8%	0.4%	5,230	83.9%	0.7%	8,268	63.6%	0.9%
Unknown	1	100.0%	0.0%	1	100.0%	0.0%	19	84.2%	8.4%	66	58.3%	6.9%

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Dialysis needed within first week excludes patients whose graft failed before the end of the week.

Table 5.14a (Continued)
Unadjusted Patient Survival, Deceased Donor non-ECD Kidney Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2007 - 2008)			(Tx 2007 - 2008)			(Tx 2003 - 2008)			(Tx 1998 - 2008)		
	N	%	Std. Err.									
Total												
All	14,326	98.5%	0.1%	14,326	96.6%	0.2%	40,616	85.0%	0.2%	69,684	64.9%	0.3%
Transplant Center State												
Alabama	248	99.2%	0.6%	248	98.4%	0.8%	651	85.2%	1.8%	1,275	63.9%	1.9%
Arkansas	140	99.3%	0.7%	140	97.9%	1.2%	407	86.5%	2.2%	598	58.1%	3.8%
Arizona	255	99.6%	0.4%	255	97.6%	0.9%	641	89.0%	1.7%	1,048	69.7%	2.3%
California	1,689	99.1%	0.2%	1,689	97.7%	0.4%	4,938	87.2%	0.6%	8,224	70.4%	0.8%
Colorado	146	99.3%	0.7%	146	96.6%	1.5%	528	88.7%	1.7%	885	65.3%	2.7%
Connecticut	108	99.1%	0.9%	108	99.1%	0.9%	336	87.7%	2.3%	564	67.2%	3.2%
District of Columbia	131	99.2%	0.8%	131	98.5%	1.1%	433	85.0%	2.2%	708	66.1%	2.9%
Delaware	17	100.0%	0.0%	17	100.0%	0.0%	24	100.0%	0.0%	30	100.0%	0.0%
Florida	1,036	98.7%	0.3%	1,036	97.7%	0.5%	2,789	84.9%	0.9%	4,772	64.3%	1.1%
Georgia	362	97.8%	0.8%	362	95.9%	1.0%	1,075	85.9%	1.4%	1,914	67.1%	1.7%
Hawaii	48	93.8%	3.5%	48	91.7%	4.0%	149	89.4%	2.9%	253	76.8%	4.0%
Iowa	120	98.3%	1.2%	120	95.0%	2.0%	324	82.2%	2.8%	594	56.8%	3.2%
Illinois	449	98.4%	0.6%	449	97.3%	0.8%	1,378	86.6%	1.1%	2,510	66.4%	1.5%
Indiana	198	98.0%	1.0%	198	97.5%	1.1%	679	87.7%	1.5%	1,188	74.6%	2.0%
Kansas	140	98.6%	1.0%	140	95.7%	1.7%	362	89.9%	1.9%	602	70.5%	3.0%
Kentucky	134	100.0%	0.0%	134	99.3%	0.7%	447	83.3%	2.2%	829	58.9%	2.6%
Louisiana	264	98.5%	0.8%	264	96.6%	1.1%	865	82.8%	1.6%	1,441	62.4%	2.1%
Massachusetts	255	96.1%	1.2%	255	94.9%	1.4%	773	86.9%	1.5%	1,405	65.4%	2.0%
Maryland	282	97.9%	0.9%	282	95.7%	1.2%	762	81.9%	1.8%	1,599	55.0%	1.8%
Maine	43	100.0%	0.0%	43	95.3%	3.2%	154	84.4%	3.4%	270	59.1%	5.0%
Michigan	525	97.3%	0.7%	525	95.0%	0.9%	1,388	82.2%	1.3%	2,357	61.6%	1.6%
Minnesota	181	100.0%	0.0%	181	98.9%	0.8%	583	85.4%	1.8%	1,046	65.4%	2.2%
Missouri	419	100.0%	0.0%	419	98.1%	0.7%	1,124	86.5%	1.4%	1,938	65.7%	1.7%
Mississippi	156	97.4%	1.3%	156	97.4%	1.3%	233	88.6%	3.2%	364	78.4%	3.4%
North Carolina	511	98.2%	0.6%	511	96.9%	0.8%	1,272	85.3%	1.4%	1,944	68.2%	1.8%
North Dakota	33	97.0%	3.0%	33	93.9%	4.2%	93	70.6%	6.4%	170	43.9%	6.6%
Nebraska	65	100.0%	0.0%	65	95.4%	2.6%	211	81.1%	3.3%	371	68.1%	3.7%
New Hampshire	28	100.0%	0.0%	28	96.4%	3.5%	88	83.2%	4.5%	139	79.0%	4.3%
New Jersey	335	95.2%	1.2%	335	92.2%	1.5%	845	80.0%	1.7%	1,429	59.2%	2.1%
New Mexico	82	98.8%	1.2%	82	97.6%	1.7%	223	94.0%	2.1%	363	70.7%	4.1%
Nevada	78	100.0%	0.0%	78	94.9%	2.5%	245	77.3%	3.4%	418	52.0%	3.8%
New York	876	98.5%	0.4%	876	95.3%	0.7%	2,315	84.1%	1.0%	3,824	65.4%	1.2%
Ohio	527	98.7%	0.5%	527	96.0%	0.9%	1,388	79.3%	1.4%	2,661	59.0%	1.4%
Oklahoma	135	100.0%	0.0%	135	97.0%	1.5%	424	78.8%	2.5%	776	59.3%	2.6%
Oregon	155	99.4%	0.6%	155	97.4%	1.3%	489	86.3%	2.1%	828	67.1%	2.6%

(Continued)

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Dialysis needed within first week excludes patients whose graft failed before the end of the week.

Table 5.14a (Continued)
Unadjusted Patient Survival, Deceased Donor non-ECD Kidney Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2007 - 2008)			(Tx 2007 - 2008)			(Tx 2003 - 2008)			(Tx 1998 - 2008)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Transplant Center State												
Pennsylvania	938	98.3%	0.4%	938	96.2%	0.6%	2,826	82.1%	0.9%	4,811	62.6%	1.1%
Puerto Rico	116	99.1%	0.9%	116	99.1%	0.9%	339	91.5%	1.9%	511	67.4%	3.9%
Rhode Island	25	100.0%	0.0%	25	96.0%	3.9%	123	81.5%	4.1%	241	51.7%	5.1%
South Carolina	210	100.0%	0.0%	210	97.1%	1.1%	582	88.4%	1.7%	1,013	63.4%	2.5%
South Dakota	33	100.0%	0.0%	33	97.0%	3.0%	92	88.4%	4.5%	178	70.5%	4.6%
Tennessee	391	98.5%	0.6%	391	95.9%	1.0%	1,134	84.3%	1.3%	1,899	58.6%	1.9%
Texas	1,229	98.2%	0.4%	1,229	96.5%	0.5%	3,502	85.8%	0.7%	6,017	66.4%	1.0%
Utah	105	100.0%	0.0%	105	99.0%	0.9%	310	92.1%	1.9%	553	73.3%	2.9%
Virginia	340	96.8%	1.0%	340	95.0%	1.2%	894	83.5%	1.7%	1,488	64.4%	1.9%
Vermont	30	100.0%	0.0%	30	93.3%	4.6%	85	79.5%	6.5%	116	75.3%	6.3%
Washington	314	99.4%	0.4%	314	98.7%	0.6%	916	89.2%	1.3%	1,495	65.4%	2.1%
Wisconsin	362	98.1%	0.7%	362	95.6%	1.1%	991	85.2%	1.4%	1,662	64.3%	1.9%
West Virginia	62	96.8%	2.2%	62	95.2%	2.7%	186	75.5%	3.9%	363	53.0%	3.8%

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Dialysis needed within first week excludes patients whose graft failed before the end of the week.