

Table 5.15b
Unadjusted Patient Survival by Year of Transplant at 3 Months, 1 Year,
3 Years, 5 Years and 10 Years
Deceased Donor ECD Kidney Transplants

Trans- # plant Year	3 Months		1 Year		3 Years		5 Years		10 Years		
	Surv.	Std. Err.	Surv.	Std. Err.	Surv.	Std. Err.	Surv.	Std. Err.	Surv.	Std. Err.	
Trans- # plant Year											
1987	37	83.8%	6.1%	75.7%	7.1%	75.7%	7.1%	64.9%	7.8%	43.2%	8.1%
1988	149	97.3%	1.3%	93.3%	2.0%	75.8%	3.5%	60.4%	4.0%	43.6%	4.1%
1989	192	95.8%	1.4%	90.1%	2.2%	82.3%	2.8%	71.4%	3.3%	49.0%	3.6%
1990	286	95.8%	1.2%	88.1%	1.9%	79.0%	2.4%	72.4%	2.6%	47.9%	3.0%
1991	348	96.3%	1.0%	92.5%	1.4%	83.3%	2.0%	72.7%	2.4%	47.1%	2.7%
1992	434	95.9%	1.0%	91.5%	1.3%	81.1%	1.9%	71.0%	2.2%	46.5%	2.4%
1993	471	94.3%	1.1%	89.2%	1.4%	79.4%	1.9%	67.1%	2.2%	45.2%	2.3%
1994	738	97.7%	0.6%	92.8%	1.0%	83.6%	1.4%	73.2%	1.6%	48.0%	1.8%
1995	883	96.1%	0.6%	92.1%	0.9%	83.5%	1.3%	72.9%	1.5%	48.2%	1.7%
1996	971	96.2%	0.6%	91.9%	0.9%	83.1%	1.2%	72.5%	1.4%	48.4%	1.6%
1997	1,020	95.3%	0.7%	88.9%	1.0%	78.9%	1.3%	68.2%	1.5%	44.6%	1.6%
1998	1,084	96.8%	0.5%	90.4%	0.9%	79.5%	1.2%	69.9%	1.4%	45.9%	1.5%
1999	1,101	95.4%	0.6%	90.5%	0.9%	80.8%	1.2%	69.8%	1.4%	47.1%	1.5%
2000	1,044	96.0%	0.6%	90.1%	0.9%	81.4%	1.2%	69.5%	1.4%	46.1%	1.7%
2001	1,084	95.7%	0.6%	89.0%	0.9%	79.6%	1.2%	67.0%	1.4%	+	+
2002	1,184	95.4%	0.6%	90.8%	0.8%	81.4%	1.1%	71.2%	1.3%	+	+
2003	1,324	96.2%	0.5%	90.1%	0.8%	80.3%	1.1%	70.9%	1.2%	+	+
2004	1,399	96.6%	0.5%	90.6%	0.8%	79.6%	1.1%	69.0%	1.2%	+	+
2005	1,592	96.6%	0.5%	90.3%	0.7%	81.7%	1.0%	71.8%	1.3%	+	+
2006	1,624	96.1%	0.5%	91.3%	0.7%	82.4%	0.9%	+	+	+	+
2007	1,666	97.5%	0.4%	93.3%	0.6%	85.7%	1.0%	+	+	+	+
2008	1,768	97.7%	0.4%	93.3%	0.6%	+	+	+	+	+	+

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

(+) = Values not determined due to insufficient follow-up.

Multi-organ transplants are excluded for kidney, pancreas, heart, lung, and liver survival rates.

Intestine survival includes intestine only and intestine with liver and/or pancreas transplants.

Living donor transplants excluded unless explicitly listed. Heterotopic heart and liver transplants excluded.

Patient survival follows patients from first transplant of this type until death. Graft survival follows individual transplants until graft failure. Counts for patient and graft survival are different because a patient may have more than one transplant for a type organ.

See Technical Notes for complete discussion of patient and graft survival methods. See Technical Notes for details of patient and graft survival methods.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDs.