

**Table 9.6f**  
**Immunosuppression Use for Maintenance by Regimen**  
**One Year Following Transplantation, 1999 to 2008**  
**Recipients with Liver Transplants**

	Transplant Year									
	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
<b>Transplants</b>	4,755	5,001	5,195	5,332	5,673	6,171	6,444	6,651	6,494	6,319
<b>Functioning Graft at 1 Year PostTx</b>	3,658	3,900	4,095	4,307	4,579	5,013	5,124	5,447	5,397	5,254
<b>With Maintenance Use Recorded at 1 Year PostTx</b>	3,411	3,715	4,027	4,210	4,441	4,899	4,999	5,266	5,122	4,898
CyA	4.2%	3.1%	2.5%	2.5%	1.9%	1.9%	1.7%	1.9%	2.1%	1.9%
Tac	20.5%	23.6%	28.9%	31.3%	30.1%	28.3%	28.6%	26.6%	28.5%	28.3%
--- + MMF/MPA	0.1%	0.0%	0.1%	0.2%	0.4%	0.3%	0.3%	0.4%	0.4%	0.8%
CyA + MMF/MPA	1.6%	1.3%	1.4%	1.6%	1.9%	2.4%	2.9%	3.0%	2.9%	3.3%
Tac + MMF/MPA	5.5%	7.1%	7.3%	10.1%	13.7%	15.2%	18.1%	22.1%	25.1%	25.9%
--- + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
CyA + OtherAntimet	0.2%	0.1%	0.1%	0.0%	0.1%	0.2%	0.0%	0.0%	0.1%	0.0%
Tac + OtherAntimet	0.3%	0.5%	0.2%	0.5%	0.3%	0.6%	0.1%	0.2%	0.3%	0.5%
--- + MMF/MPA + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
CyA + MMF/MPA + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Tac + MMF/MPA + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
--- + Siro/Evero	0.2%	0.8%	1.0%	1.7%	1.5%	2.0%	2.0%	1.9%	1.7%	1.6%
CyA + Siro/Evero	0.1%	0.9%	0.5%	0.3%	0.4%	0.3%	0.5%	0.3%	0.3%	0.3%
Tac + Siro/Evero	0.4%	2.4%	2.5%	2.3%	2.4%	2.9%	2.1%	1.2%	1.2%	1.2%
--- + MMF/MPA + Siro/Evero	0.0%	0.2%	0.4%	1.0%	1.1%	1.3%	2.4%	2.2%	1.8%	1.2%
CyA + MMF/MPA + Siro/Evero	0.1%	0.0%	0.0%	0.0%	0.0%	0.1%	0.1%	0.0%	0.0%	0.0%
Tac + MMF/MPA + Siro/Evero	0.0%	0.2%	0.1%	0.2%	0.3%	0.2%	0.2%	0.2%	0.3%	0.2%
--- + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.1%	0.0%	0.0%	0.0%
Tac + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
--- + Steroids	1.8%	1.6%	0.6%	0.8%	0.5%	0.4%	0.3%	0.2%	1.0%	1.5%
CyA + Steroids	10.3%	5.8%	2.9%	2.3%	2.0%	1.7%	1.2%	1.2%	1.0%	0.9%
Tac + Steroids	30.5%	26.7%	22.8%	22.3%	20.9%	18.1%	15.7%	15.2%	12.6%	11.3%
--- + MMF/MPA + Steroids	0.4%	0.3%	0.3%	0.4%	0.4%	0.4%	0.4%	0.5%	0.4%	0.6%
CyA + MMF/MPA + Steroids	5.0%	4.0%	3.2%	2.0%	2.1%	2.0%	1.7%	1.5%	1.5%	1.5%
Tac + MMF/MPA + Steroids	10.4%	10.8%	13.2%	12.6%	14.1%	15.5%	16.8%	17.4%	15.1%	15.4%
--- + OtherAntimet + Steroids	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%	0.0%
CyA + OtherAntimet + Steroids	2.5%	1.7%	0.8%	0.3%	0.2%	0.3%	0.1%	0.0%	0.1%	0.0%
Tac + OtherAntimet + Steroids	1.4%	1.1%	1.4%	1.4%	1.0%	0.5%	0.3%	0.3%	0.3%	0.2%
--- + Siro/Evero + Steroids	0.1%	0.3%	1.0%	0.9%	0.8%	1.6%	1.3%	1.0%	0.9%	0.9%
CyA + Siro/Evero + Steroids	0.3%	0.3%	0.4%	0.3%	0.3%	0.1%	0.1%	0.0%	0.1%	0.2%
Tac + Siro/Evero + Steroids	0.6%	3.0%	2.3%	2.1%	1.5%	1.5%	1.2%	0.9%	0.8%	0.9%
Other Regimen	3.6%	4.3%	5.8%	2.7%	2.1%	1.8%	1.6%	1.7%	1.5%	1.3%

Source: OPTN/SRTR Data as of October 1, 2010.

Notes:

Percentages are calculated based on the total number of patients with maintenance use at one year following transplantation. They sum to 100 percent for each column.

---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.

MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.

"Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.